

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁶ : C12P 21/08, C07K 16/40, G01N 33/543, C12N 9/26, 15/52, A61K 38/46, 39/395, 31/70	A1	(11) International Publication Number: WO 98/16655 (43) International Publication Date: 23 April 1998 (23.04.98)
(21) International Application Number: PCT/US97/18089 (22) International Filing Date: 7 October 1997 (07.10.97) (30) Priority Data: 08/733,360 17 October 1996 (17.10.96) US 08/916,935 21 August 1997 (21.08.97) US (71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 22nd floor, 300 Lakeside Drive, Oakland, CA 94612-3550 (US). (72) Inventors: STERN, Robert; 598 Belvedere Street, San Francisco, CA 94117 (US). FROST, Gregory, I.; 3575 Fillmore Street, San Francisco, CA 94123 (US). CSOKA, Anthony; 598 Belvedere Street, San Francisco, CA 94117 (US). WONG, Tim, M.; 2340 Wawona Street, San Francisco, CA 94116 (US). (74) Agent: BERLINER, Robert; Robbins, Berliner & Carson, LLP, 5th floor, 201 N. Figueroa Street, Los Angeles, CA 90012-2628 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: HUMAN PLASMA HYALURONIDASE		
(57) Abstract <p>The invention is based on the discovery of methods for purification of an acid active hyaluronidase found in human plasma (hpHase), including both biochemical and immunoaffinity purification methods. The method of immunoaffinity purification of the invention is based on the discovery of a method for identifying antibodies that specifically bind native hpHase (anti-native hpHase antibodies), and anti-native hpHase antibodies identified by this screening method. The invention also features an assay for sensitive detection of Hase activity using biotinylated hyaluronic acid (bHA). Purification and characterization of hpHase lead to the inventors' additional discovery that hpHase is encoded by the LuCa-1 gene, which gene is present in the human chromosome at 3p21.3, a region associated with tumor suppression. The invention additionally features methods of treating tumor-bearing patients by administration of hpHase and/or transformation of cells with hpHase-encoding DNA.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HUMAN PLASMA HYALURONIDASE

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of U.S. application serial no. 08/733,360, filed October 17, 1996, which application is hereby incorporated by reference.

STATEMENT REGARDING FEDERALLY SPONSORED
RESEARCH OR DEVELOPMENT

This invention was made with government support under grant nos. CA44768, CA58207, and GM46765, awarded by the National Institutes of Health. The government may have certain rights in this invention.

FIELD OF THE INVENTION

This invention relates generally to the field of β -1,4-endoglycosidases, particularly hyaluronidases.

BACKGROUND OF THE INVENTION

Hyaluronidases (HAs; E.C. 3.1.25) are a group of neutral- and acid-active enzymes found throughout the animal kingdom in organisms as diverse as microbes (e.g., *Streptococcus pyogenes*, *Treponema palladium*, and nematodes), bees, wasps, hornet, spiders, scorpions, fish, snakes, lizards, and mammals. Hyaluronidases degrade hyaluronan (HA; also known as hyaluronic acid) and, to a lesser extent, chondroitin sulfates (for a review, see Kreil et al. 1995 *Protein Sci.* 4:1666-9). Vertebrate hyaluronidases are separated into two general classes: 1) the neutral hyaluronidases, such as the predominantly sperm-associated protein PH20 (Liu et al. 1996 *Proc. Natl. Acad. Sci. USA* 93:7832-7; Primakoff et al. 1985 *J. Cell Biol.* 101:2239-44; Lin et al. 1993 *Proc. Natl. Acad. Sci. USA* 90:10071-5); and 2) the acid-active hyaluronidases, which have a distinct pH optimum between pH 3.5 to 4.0 and have been described in extracts of liver (Fischer-Szafarz et al. 1995 *Acta Biochim Pol.* 42:31-3), kidney (Komender et al. 1973 *Bull. Acad. Pol.*

Sci. [Biol.] 21:637-41), lung (Thet et al. 1983 *Biochem. Biophys. Res. Commun.* 117:71-7), brain (Margolis et al. 1972 *J. Neurochem.* 19:2325-32), skin (Cashman et al. 1969 *Arch. Biochem. Biophys.* 135:387-95), placenta, macrophages, fibroblasts (Lien et al. 1990 *Biochim Biophys. Acta* 1034:318-25; Ruggiero et al. 5 1987 *J. Dent. Res.* 66:1283-7), urine (Fischer-Szafarz et al. *supra*) and human plasma (De Salegui et al. 1967 *Arch. Biochem. Biophys.* 120:60-67). Acid-active hyaluronidase activity has also been described in the sera of mammals, though some species exhibit no detectable activity at all (Fischer-Szafarz et al. 1990 *Biol. Cell* 68:95-100; De Salegui et al. 1967 *supra*).

10 Hyaluronan, the main substrate for hyaluronidase, is a repeating disaccharide of $[\text{GlcNAc}\beta 1\text{-4GlcUA}\beta 1\text{-3}]_n$ that exists *in vivo* as a high molecular weight linear polysaccharide. Degradation of hyaluronan by hyaluronidase is accomplished by either cleavage at β -N-acetyl-hexosamine-[1 \rightarrow 4]-glycosidic bonds or cleavage at β -gluconate-[1 \rightarrow 3]-N-acetylglucosamine bonds.

15 Hyaluronan is found in mammals predominantly in connective tissues, skin, cartilage, and in synovial fluid. Hyaluronan is also the main constituent of the vitreous of the eye. In connective tissue, the water of hydration associated with hyaluronan creates spaces between tissues, thus creating an environment conducive to cell movement and proliferation. Hyaluronan plays a key role in biological 20 phenomena associated with cell motility including rapid development, regeneration, repair, embryogenesis, embryological development, wound healing, angiogenesis, and tumorigenesis (Toole 1991 *Cell Biol. Extracell. Matrix*, Hay (ed), Plenum Press, New York, 1384-1386; Bertrand et al. 1992 *Int. J. Cancer* 52:1-6; Knudson et al, 1993 *FASEB J.* 7:1233-1241). In addition, hyaluronan levels 25 correlate with tumor aggressiveness (Ozello et al. 1960 *Cancer Res.* 20:600-604; Takeuchi et al. 1976, *Cancer Res.* 36:2133-2139; Kimata et al. 1983 *Cancer Res.* 43:1347-1354).

Hyaluronidase is useful as a therapeutic in the treatment of diseases associated with excess hyaluronan and to enhance circulation of physiological fluids 30 and/or therapeutic agents at the site of administration. For example, hyaluronidase has been used to reduce intraocular pressure in the eyes of glaucoma patients

through degradation of hyaluronan within the vitreous humor (USPN 4,820,516, issued April 11, 1989). Hyaluronidase has also been used in cancer therapy as a "spreading agent" to enhance the activity of chemotherapeutics and/or the accessibility of tumors to chemotherapeutics (Schüller et al., 1991, *Proc. Amer. Assoc. Cancer Res.* 32:173, abstract no. 1034; Czejka et al., 1990, *Pharmazie* 45:H.9) and has been used in combination with other chemotherapeutic agents in the treatment of a variety of cancers including urinary bladder cancer (Horn et al., 1985, *J. Surg. Oncol.*, 28:304-307), squamous cell carcinoma (Kohno et al., 94, *J. Cancer Res. Oncol.*, 120:293-297), breast cancer (Beckenlehner et al., 1992, *J. Cancer Res. Oncol.* 118:591-596), and gastrointestinal cancer (Scheithauer et al., 1988, *Anticancer Res.* 8:391-396). Administration of hyaluronidase also induces responsiveness of previously chemotherapy-resistant tumors of the pancreas, stomach, colon, ovaries, and breast (Baumgartner et al., 1988, *Reg. Cancer Treat.* 1:55-58; Zänker et al., 1986, *Proc. Amer. Assoc. Cancer Res.* 27:390). Serum hyaluronidase prevents growth of tumors transplanted into mice (De Maeyer et al., 1992, *Int. J. Cancer* 51:657-660), while injection of hyaluronidase inhibits tumor formation caused by exposure to carcinogens (Pawlowski et al., 1979, *Int. J. Cancer* 23:105-109; Haberman et al., 1981, *Proceedings of the 17th Annual Meeting of the American Society of Clinical Oncology*, Washington, D.C., 22:105, abstract no. 415). Intravenous or intramuscular injection of hyaluronidase is effective in the treatment of brain cancer (gliomas) (PCT published application no. WO88/02261, published April 7, 1988).

Hyaluronidase expression, and levels of hyaluron, have been associated with tumor development and progression. Levels of a secreted neutral hyaluronidase activity in carcinomas derived from ovary (Miura et al. 1995 *Anal. Biochem.* 225:333-40), prostate (Lokeshwar et al. 1996 *Cancer Res* 56:651-7), brain, melanocyte, and colon (Liu et al. 1996 *Proc. Natl. Acad. Sci. USA* 93:7832-7837) are higher than in normal tissue. This secreted neutral hyaluronidase activity appears similar or identical to the neutral hyaluronidase activity of the sperm hyaluronidase PH20. In contrast to neutral activity, the acid active serum hyaluronidase activity is significantly decreased in metastatic carcinomas of the

lung, breast, and colon (Northrup et al. 1973 *Clin. Biochem.* 6:220-8; Kolarova et al. 1970 *Neoplasma* 17:641-8). Further, mice having an allele of the hyal-1 locus that is associated with lower levels of serum hyaluronidase activity exhibit faster rates of growth of transplanted tumors than mice having an hyal-1 allele that is associated with 3-fold higher hyaluronidase activity levels (Fischer-Szafarz et al. 1989 *Somat. Cell. Mol. Genet.* 15:79-83; De Maeyer et al. *supra*).

At present, the only hyaluronidase activity available for clinical use is a hyaluronidase isolated from a testicular extract from cattle (WYDASE®, Wyeth-Ayerst). The bovine extract is not optimum not only because of its non-human source, but also because the extract contains multiple types of hyaluronidases and other as yet undefined components. While the human serum acid-active hyaluronidase activity would be preferred for administration, this hyaluronidase has not been previously isolated or purified. Although previous studies were able to determine that the serum acid-active hyaluronidase activity is not a component of platelets since hyaluronidase activity levels in plasma are comparable to those found in serum (De Saegui et al. 1967 *supra*), attempts to isolate this acid active hyaluronidase activity from human serum have met with limited success due in part to the stability of the purified activity and the inability to obtain an adequately high specific activity. Immunopurification attempts have been hindered by the inability to identify and isolate antibodies that specifically bind the activity in its native form in plasma. Monoclonal antibodies identified by conventional ELISA techniques bind denatured human plasma hyaluronidase in the ELISA screening assay do not bind the native, folded protein (Harrison et al. 1988 *J Reprod Fertil* 82:777-85).

Conventional methods for hyaluronidase activity include ELISA-like assays (Delpech et al. 1987 *J. Immunol. Methods* 104:223-9; Stern et al. 1992 *Matrix* 12:397-403; Afify et al. 1993 *Arch. Biochem. Biophys.* 305:434-41; Reissig et al., 1955, *J. Biol. Chem.* 217:956-966) in which a sample containing hyaluronidase is applied to the well of a microtiter dish having hyaluronan or hyaluronectin non-covalently bound to its surface. Hase present in the sample degrades the HA substrate. The plates are then washed, and Hase activity is detected by examining

the plates for remaining HA substrate.

Hyaluronidase activity can also be detected by substrate gel zymography (Guentenhoner et al., 1992, *Matrix* 12:388-396). In this assay a sample is applied to a SDS-PAGE gel containing hyaluronan and the proteins in the sample separated
5 by electrophoresis. The gel is then incubated in an enzyme assay buffer and subsequently stained to detect the hyaluronan in the gel. Hyaluronidase activity is visualized as a cleared zone in the substrate gel.

These conventional methods for detecting hyaluronidase activity are hampered by both the difficulties in producing a detectably-labeled hyaluronic acid
10 substrate and the technical difficulties in achieving rapid, sensitive, and reproducible detection of hyaluronidase activity. For example, biotin labeling of hyaluronic acid for use in ELISA-like assays has proved reticent to biotinylation since HA contains no free amine groups, the moiety with which activated biotin covalently binds. Prior attempts to solve this problem have focused on use of a
15 biotinylated-HA binding aggrecan peptide from bovine nasal cartilage (Levvy et al. 1966 *Method Enzymol.* 8:571-584), which requires tedious, time-consuming steps.

Furthermore, conventional hyaluronidase assays use assay plates having HA substrate non-covalently bound to the plate surface, which can lead to both false
20 positive and false negative results. Because the HA substrate is non-covalently bound to the plate surface, the washing step following exposure of the plates to the Hase-containing sample often results in non-specific removal of non-degraded HA substrate on the plate. Thus, the sensitivity of the conventional Hase assay is compromised. Hase activities using gel zymography avoid the problem associated
25 with ELISA-like assays, but are time-consuming (e.g., the test sample and the HA-containing gel are normally incubated for about 18 hr to 24 hr) and can result in artifacts if the gel is improperly loaded with excess protein sample. Moreover, analyses of crude preparations is impossible by gel zymography.

Thus, despite the presence of a desirable acid active plasma hyaluronidase
30 activity, and human blood product companies' economic motivation to obtain any and all useful components from a resource as precious and scarce as human blood,

the human plasma fractions containing this acid active hyaluronidase activity are discarded for want of an acceptable method for its isolation and purification.

Given the value of hyaluronidases in chemotherapy, there is a need in the field for a method of identifying and isolating the polypeptide associated with the acid active hyaluronidase activity in human serum.

SUMMARY OF THE INVENTION

The invention is based on the discovery of methods for purification of an acid active hyaluronidase found in human plasma (hpHase), which include both biochemical and immunoaffinity purification methods. The method of immunoaffinity purification of the invention is based on the discovery of a method for identifying antibodies that specifically bind native hpHase (anti-native hpHase antibodies), and anti-native hpHase antibodies identified by this screening method. The invention also features an assay for sensitive detection of Hase activity using biotinylated hyaluronic acid (bHA). Purification and characterization of hpHase lead to the inventors' additional discovery that hpHase is encoded by the LuCa-1 gene, which is located in the human chromosome at 3p21.3, a region associated with tumor suppression.

Thus, in one aspect the invention features an assay device for detection of hyaluronidase activity comprising an insoluble support and biotinylated hyaluronic acid (bHA) covalently bound to the support. Preferably, the bHA is prepared in a one-step reaction of hyaluronic acid, 1-ethyl-dimethylaminopropyl carbodiimide (EDC), and biotin hydrazide. In preferred embodiments, the bHA comprises at least one biotin moiety per every 100 disaccharide units in the hyaluronic acid moiety, and the bHA is covalently bound to the support by at least one covalent bond per every 50 disaccharide units in the hyaluronic acid moiety.

In another aspect the invention features a method of purifying a native acid active hyaluronidase (aaHase) from a sample, the method comprising the steps of: (a) dissolving a sample suspected of containing aaHase in a solution at a temperature substantially less than room temperature, the solution comprising a non-ionic detergent; (b) raising the temperature of the solution to a temperature

substantially greater than room temperature, said raising resulting in the formation of a detergent-rich phase comprising aaHase and a detergent-poor phase; and (c) isolating aaHase from the detergent-rich phase. Preferably the aaHase is hpHase and steps (a)-(c) are repeated twice.

5 In yet another aspect of the invention, the invention features a method for screening candidate antibodies for binding to a native aaHase comprising the steps of: (a) incubating a candidate antibody with a sample comprising native aaHase, said incubating being for a time sufficient for formation of antibody-aaHase complexes; (b) contacting the sample with an insoluble support having anti-
10 antibody and detectably-labeled hyaluronic acid bound thereto for a time sufficient for formation of anti-antibody-candidate antibody-hpHase complexes; and (c) exposing the sample in contact with the support to an acidic pH of about 3.4 to 3.7, thereby allowing hpHase in the antibody-aaHase complex to degrade the
15 degradation comprise an anti-aaHase antibody. In related aspects, the invention features anti-native hpHase antibodies, hybridoma cell lines secreting such antibodies, and assay devices for immunopurification and/or detection of hpHase comprising anti-hpHase antibodies bound thereto.

 In still another aspect, the invention features a method of purifying hpHase
20 from a sample, the method comprising contacting a sample comprising hpHase with an anti-hpHase antibody. In preferred embodiments, the sample is human blood, serum, plasma, or urine or the hpHase is a recombinant hpHase.

 In another aspect, the invention features substantially purified native hpHase characterized by a fatty acid moiety that is resistant to cleavage by
25 phospholipase C, phospholipase D, and N-glycosidase-F. In related aspects the invention features formulations comprising native hpHase, particularly liposome formulations.

 In yet another aspect the invention features methods for recombinantly expressing hpHase. The recombinant expression system of the invention provides
30 high levels of secreted hpHase suitable for commercial production of hpHase and for production of hpHase for use in therapeutic applications. In related aspects

the invention features recombinant hpHase and methods of making recombinant hpHase.

In additional related aspects, the invention features methods of treating patients having or susceptible to a condition for which administration of hyaluronidase is desirable. In specific embodiments, the invention features administration of human plasma hyaluronidase to a patient having or susceptible to cancer associated with defective hpHase expression comprising administering hpHase polypeptide to the patient in an amount effective to suppress tumor growth. In further embodiments, the invention features methods of treating a patient after suffering a myocardial infarction or a patient suffering from a lysosomal storage disease.

Another aspect of the invention features a method of treating a patient having or susceptible to cancer associated with a defective LuCa-1 gene, the method comprising introducing into a cell of the patient a construct comprising a nucleotide sequence encoding a human plasma hyaluronidase polypeptide and a eukaryotic promoting sequence operably linked thereto, resulting in the genetic transformation of the cell so that the nucleotide sequence expresses hpHase.

The invention also features a method for identifying a patient having or susceptible to a condition associated with defective human plasma hyaluronidase activity comprising the steps of contacting a plasma sample from the patient with an anti-native human plasma hyaluronidase antibody and detecting complexes of the enzyme and the antibody.

A primary object of the invention is to provide a purified hyaluronidase, hpHase, which can be used in a variety of clinical therapies including cancer therapy, particularly cancers associated with a defect in the tumor suppressor gene LuCa-1.

An advantage of the present invention is that purified hpHase is more appropriate for therapeutic uses than the presently available commercial formulations of hyaluronidase which are from a non-human source, which contain two hyaluronidases (rather than one), and which, as determined by SDS-PAGE analysis, is a very crude mixture that contains various proteins, including several

unidentified proteins and proteins having various biological activities including anticoagulant activities (Doctor et al., Thrombosis Res. 30:565-571). Purified hpHase provides a "clean" source of hyaluronidase, is less likely to induce some of the side effects associated with the presently available commercial formulation, and allows better control of the level of activity associated with specific dosages.

Another advantage of the invention is that hpHase can be purified from a lipid fraction of plasma that, according to present commercial practices, is otherwise discarded.

An advantage of the Hase assay of the invention is that it provides for an assay for Hase activity that is at least 1,000-fold more sensitive than conventional ELISA-like assays. In addition, the invention provides a biotinylated hyaluronic acid substrate that can be easily and efficiently prepared.

An advantage of the anti-native aaHase assay of the invention is that the assay provides a rapid screen for antibodies that bind native aaHase.

Another advantage of the invention is that specific detection of hpHase allows a specific means of measuring LuCa-1 expression and correlation of levels of hpHase in serum or urine of a patient with susceptibility to or presence of a disease associated with a LuCa-1 defect (e.g., cancer, e.g., small cell lung cell carcinoma).

These and other objects, advantages and features of the present invention will become apparent to those persons skilled in the art upon reading the details of the invention as more fully set forth below.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a schematic showing the chemical and physical configuration of a surface for covalent attachment of biotinylated HA.

Fig. 2 is a schematic showing the chemical structure of biotinylated HA.

Fig. 3 is a schematic showing the steps in the anti-native acid active Hase antibody assay of the invention.

Fig. 4 is a graph showing hyaluronidase activity associated with supernatants of human plasma after immunoprecipitation with varying

concentrations of anti-native hpHase 17E9 antibody (open circles) or bound control antibody (open squares).

Fig. 5 is a schematic showing the alignment of the predicted amino acid sequence of LuCa-1 (SEQ ID NO:3) with the amino acid sequence of the neutral
5 Hase PH20 (SEQ ID NO:11). Identical amino acid residues are shaded.

Fig. 6 is a hydropathy plot of the LuCa-1/hpHase.

Fig. 7A is a graph showing the acid active Hase activity of detergent extracts of LuCa-1-expressing cos-7 cells and conditioned medium (shaded
10 columns) and of mock-transfected cells and conditioned medium (unshaded columns).

Fig. 7B is a graph showing the acid active Hase activity of detergent extracts of LuCa-1 expressing Hek 293 cells and conditioned medium (shaded columns) and of mock-transfected cells and conditioned medium (unshaded columns).

15 Fig. 8 is a graph showing the pH optima of the Hase activity of hpHase purified from human plasma (open squares and solid line), recombinant LuCa-1 expressed in cos-7 cells (closed squares with dashed line), and Type VI-S testicular hyaluronidase (closed circles with dashed line).

Fig. 9 is a graph showing hpHase expression in the cell layer and
20 conditioned media of normal human keratinocytes and several carcinoma cell lines.

Fig. 10 is a graph showing the effects of hpHase peritumoral injection upon growth of HSC-3 head and neck squamous cell carcinoma cells in an animal model.

Fig. 11 is a graph showing the effects of hpHase expression in transformed
25 HSC-3 cells upon tumor growth in an animal model.

Fig. 12 is a graph showing the effects of hpHase expression in transformed HSC-3 cells upon cachexia in an animal model.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Before the present purified hyaluronidase and DNA encoding same are described, it is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors and reagents described as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a transformed cell containing DNA encoding a hyaluronidase" includes a plurality of such cells and reference to "the transformation vector" includes reference to one or more transformation vectors and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs. Although any methods, devices and materials similar or equivalent to those described herein can be used in the practice or testing of the invention, the preferred methods, devices and materials are now described.

All publications mentioned herein are incorporated herein by reference for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are described in the publications which might be used in connection with the presently described invention.

Definitions

By "acid active hyaluronidase" or "aaHase" is meant a hyaluronidase having β -1,4-endoglycosidase activity in the cleavage of hyaluronan and a pH optimum of Hase activity at about pH 3.7. aaHase as used herein encompasses human plasma hyaluronidase.

By "human plasma hyaluronidase," "human plasma acid active hyaluronidase," and "hpHase" is meant a hyaluronidase naturally found in human

plasma and having the following characteristics: 1) β -1,4-endoglycosidase activity in the cleavage of hyaluron; 2) a pH optimum of Hase activity at about pH 3.7; 3) a molecular weight of about 57 kDa as determined by 12.5% SDS-PAGE non-reducing gel electrophoresis; 4) a specific enzymatic activity of about 2×10^5 to 8×10^5 turbidity reducing units (TRU)/mg protein following purification; 5) an isoelectric point, as determined by elution in chromatofocusing on Mono-P f.p.l.c., of pH 6.5; 6) partitioning into the Triton X-114 detergent-rich phase upon temperature-induced detergent phase extraction; 7) a fatty acid post-translational modification (e.g., a lipid anchor) that is resistant to cleavage by phospholipase C, phospholipase D, and N-glycosidase-F; 8) at least two N-linked glycosylation sites; and 9) has the amino acid sequence:

MAGHLLPICALFLTLLDMAQGRGPLLPNRPFTTVWNANTQWCLERHGV
 DVDVSVFDVVANPGQTFRGPDMTIFYSSQLGTYPYYTPTGEPVFGGLPQN
 ASLIAHLARTFQDILAAIPAPDFSGLAVIDWEAWRPRWAFNWDTKDIYRQ
 15 RSRALVQAQHPDWPAPQVEAVAQDQFQGAARAWMAGTLQLGGALRPRG
 LWGFYGFPCYNYDFLSPNYTGQCPSGIRAQNDQLGWLWGQSRALYPSI
 YMPAVLEGTGKSQMYVQHRVAEAFRVAVAAGDPNLPVLPYVQIFYDTTN
 HFLPLDELEHSLGESAAQGAAGVVLWVSWENTRTKESCQAIKEYMDTTL
 GPFILNVTSGALLCSQALCSGHGRCVRRTSHPKALLLNPAFSIQLTPGGG
 20 PLSLRGALSLEDQAQMAVEFKCRCYPGWQAPWCERKSMW (SEQ ID NO:1)

where MAGHLLPICALFLTLLDMAQG (SEQ ID NO:2) is a signal sequence cleaved during post-translational modification. "hpHase" as used herein is meant to encompass hpHase polypeptides having the amino acid sequence of naturally-occurring hpHase, as well as hpHase polypeptides that are modified relative to the naturally-occurring amino acid sequence due to amino acid substitution, deletion, and/or addition (e.g., fusion proteins) and the like. Preferably, "hpHase" encompasses hpHase polypeptides that are biologically active (e.g., can bind anti-hpHase antibodies and/or exhibit hyaluronidase activity).

By "urine form of hpHase" or "urine hpHase" is meant a form of hpHase is found in human urine and characterized by: 1) β -1,4-endoglycosidase activity in the cleavage of hyaluron; 2) a pH optimum of Hase activity at about pH 3.7; 3)

a molecular weight of about 57 kDa as determined by gel zymography using 12.5% SDS-PAGE non-reducing gel electrophoresis; 4) immunoprecipitation with anti-native hpHase monoclonal antibodies specific to LuCa-1; 5) an isoelectric point, as determined by elution in chromatofocusing on Mono-P f.p.l.c., or pH 6.5; and
5 6) partitioning into the Triton X-114 detergent-rich phase upon temperature-induced detergent phase extraction.

By "native hpHase" is meant hpHase that is folded in its naturally-occurring configuration (i.e., hpHase is not denatured). Where the native hpHase does not comprise the entire amino acid sequence of naturally-occurring hpHase
10 (SEQ ID NO:1), native hpHase polypeptides are those polypeptides that, when folded, mimic a three-dimensional epitope of native, full-length hpHase such that antibodies that bind native hpHase bind to the hpHase polypeptide. "Native hpHase" encompasses both hpHase that is naturally found in human plasma, blood, serum, and urine, as well as hpHase that is recombinantly produced (e.g.,
15 by expression in a mammalian cell).

By "polypeptide" is meant any chain of amino acids, regardless of length or post-translational modification (e.g., glycosylation, phosphorylation, or fatty acid chain modification).

By a "substantially pure polypeptide" a polypeptide that has been separated
20 from components which naturally accompany it (e.g., a substantially pure hpHase polypeptide purified from human plasma is substantially free of components normally associated with human plasma). Typically, the polypeptide is substantially pure when it is at least 60%, by weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated.
25 Preferably, the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, hpHase polypeptide. A substantially pure hpHase polypeptide can be obtained, for example, by extraction from a natural source (e.g., mammalian plasma, preferably human plasma); by expression of a recombinant nucleic acid encoding hpHase polypeptide; or by chemically
30 synthesizing the protein. Purity can be measured by any appropriate method, e.g.,

chromatography, polyacrylamide gel electrophoresis, or by HPLC analysis.

A protein is substantially free of naturally associated components when it is separated from those contaminants which accompany it in its natural state. Thus, a protein which is chemically synthesized or produced in a cellular system
5 different from the cell from which it naturally originates will be substantially free from its naturally associated components. Accordingly, substantially pure polypeptides include those derived from eukaryotic organisms but synthesized in *E. coli* or other prokaryotes.

By "antibody" is meant an immunoglobulin polypeptide that is capable of
10 binding an antigen. Antibody as used herein is meant to include the entire antibody as well as any antibody fragments (e.g. $F(ab')_2$, Fab' , Fab , Fv) capable of binding the epitope, antigen or antigenic fragment of interest. Antibodies of the invention are immunoreactive or immunospecific for and therefore specifically and selectively bind to native hpHase polypeptide. Anti-hpHase antibodies are
15 preferably immunospecific (i.e., not substantially cross-reactive with related materials). Antibodies may be polyclonal or monoclonal, preferably monoclonal.

By "binds specifically" is meant high avidity and/or high affinity binding of an antibody to a specific polypeptide i.e., epitope of hpHase. Antibody binding to its epitope on this specific polypeptide is preferably stronger than binding of the
20 same antibody to any other epitope, particularly those which may be present in molecules in association with, or in the same sample, as the specific polypeptide of interest e.g., binds more strongly to hpHase than to other components in human plasma. Antibodies that bind specifically to a polypeptide of interest may be capable of binding other polypeptides at a weak, yet detectable, level (e.g., 10%
25 or less of the binding shown to the polypeptide of interest). Such weak binding, or background binding, is readily discernible from the specific antibody binding to the compound or polypeptide of interest, e.g. by use of appropriate controls. In general, antibodies of the invention which bind to native hpHase with a binding affinity of 10^7 liters/mole or more, preferably 10^8 l/mole or more, even more
30 preferably 10^9 l/mole or more, are said to bind specifically to hpHase. In general, an antibody with a binding affinity of 10^4 l/mole or less is not useful in that it will

not bind an antigen at a detectable level using conventional methodology currently used.

By "anti-native hpHase antibody" or "anti-hpHase antibody" is meant an antibody that specifically binds native (i.e., non-denatured) hpHase. Preferably, such antibodies can be used to immunopurify (e.g., by immunoprecipitation or immunoaffinity column chromatography) naturally-occurring hpHase from human plasma and/or recombinant hpHase expressed by, for example, mammalian cells.

By "operably linked" is meant that a DNA of interest (e.g., DNA encoding an hpHase polypeptide) and a regulatory sequence(s) are connected in such a way as to permit gene expression of the DNA of interest when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the regulatory sequence(s), thus facilitating production of, e.g., an hpHase polypeptide, a recombinant protein, or an RNA molecule.

By "transformation" is meant a permanent genetic change induced in a cell following incorporation of new DNA (i.e., DNA exogenous to the cell). Where the cell is a mammalian cell, the permanent genetic change is generally achieved by introduction of the DNA into the genome of the cell.

By "vector" is meant any compound, biological or chemical, which facilitates transformation of a host cell with DNA encoding an hpHase polypeptide of the invention.

By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 50%, preferably 85%, more preferably 90%, and most preferably 95% homology to a reference amino acid or nucleic acid sequence. For polypeptides, the length of comparison sequences will generally be at least 16 amino acids, preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids. For nucleic acids, the length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, and most preferably 110 nucleotides.

Sequence identity is typically measured using sequence analysis software (e.g., Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison,

WI 53705). Such software matches similar sequences by assigning degrees of homology to various substitutions, deletions, substitutions, and other modifications. Conservative substitutions typically include substitutions within the following groups: glycine alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

The terms "treatment", "treating" and the like are used herein to generally mean obtaining a desired pharmacologic and/or physiologic effect. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of a partial or complete cure for a disease and/or adverse effect attributable to the disease. "Treatment" as used herein covers any treatment of a disease in a mammal, particularly a human, and includes: (a) preventing the disease (e.g., cancer) from occurring in a subject who may be predisposed to the disease but has not yet been diagnosed as having it; (b) inhibiting the disease, i.e., arresting its development; or (c) relieving the disease, i.e., causing regression of the disease (e.g., reduction in tumor volume, slowing of cachexia). The invention is directed toward treatment of patients having or susceptible to cancer associated with a defect in the LuCa-1 gene, the gene that encodes hpHase.

By "therapeutically effective amount of a substantially pure hpHase polypeptide" is meant an amount of a substantially pure hpHase polypeptide effective to facilitate a desired therapeutic effect. The precise desired therapeutic effect will vary according to the condition to be treated. For example, the desired degradation of hyaluronan is the desired therapeutic effect where hpHase is administered to the subject in the treatment of a condition associated with excess hyaluron, undesirable cell motility (e.g., tumor cell metastasis), and/or to enhance circulation of physiological fluids at the site of administration and/or inhibit tumor growth or progression. Where hpHase is administered to treat a patient having or susceptible to cancer associated with a LuCa-1 defect, one desired therapeutic effects include, but are not necessarily limited to, an inhibition of tumor cell growth and a decrease in the tumor cell's threshold to apoptosis (i.e., increase the

cell's sensitivity to triggers to programmed cell death). Therapeutic effects of hpHase may be associated with hpHase's hyaluronidase activity, chondroitinase activity, or both.

By "LuCa-1 defect" or "hpHase defect" is meant a genetic defect in a cell
5 at chromosome position 3p.21.3 associated with a decreased level of hpHase activity relative to hpHase activity levels in normal cells and/or a decreased level of hpHase activity in the serum or plasma of the affected patient (e.g., due to decreased expression of hpHase or expression of a defective hpHase). "LuCa-1 defect" or "hpHase defect" are also meant to encompass cellular defects associated
10 with a decreased level of hpHase activity in the serum or plasma of the affected patient (e.g., a defect in transport of hpHase into the bloodstream). For example, plasma from patients have a LuCa-1 defect-associated lung cancer exhibits about 50% less hpHase activity than plasma from normal patients (i.e., patients who do not have a LuCa-1 defect-associated cancer). Normal human plasma exhibits about
15 15 rTRU/mg hpHase activity as determined using the Hase assay of the invention. The hpHase activity of plasma from LuCa-1 defect-associated lung cancer patients is about 7.5 rTRU/mg.

By "having or susceptible to a condition associated with a LuCa-1 defect" is meant to describe a patient having a heterozygous, homozygous, or epigenetic
20 defect at the LuCa-1 locus and/or some other genetic defect outside the LuCa-1 locus associated with decreased levels of hpHase (e.g., serum hpHase or urine hpHase, preferably serum hpHase) relative to hpHase levels associated with normal patients (i.e., patients having no LuCa-1 defect that results in altered hpHase levels). Exemplary patients having or susceptible to a condition associated
25 with a LuCa-1 defect are patients bearing tumor or pre-tumor cells that do not express normal levels of hpHase (e.g., a metastatic carcinoma).

The invention will now be described in further detail.

Hyaluronidase Activity Assay

The Hase assay of the invention involves labeling of HA by biotinylation of free carboxyl groups through a simple, one step reaction with 1-ethyl-dimethylaminopropyl carbodiimide (EDC, $\text{ClCH}_2\text{CH}_2\text{Cl}$, Sigma) and biotin hydrazide (Pierce). Briefly, a solution composed of dissolved HA and dissolved biotin hydrazide is combined with EDC. Preferably, biotin is present in excess relative to EDC and HA. The molar amount of EDC present is varied according to the desired number of covalent bonds to be found between the carboxyl groups of the disaccharide units of HA and the NH groups of the biotin moieties (i.e., the desired number of biotin moieties relative to the number of HA disaccharide units). Preferably, the molar ratio of HA disaccharide units to EDC is 85:1, while the molar ratio of biotin hydrazide to EDC is 38:1. An exemplary chemical structure of the resulting HA-bHA compound is shown in Fig. 1.

After the HA-EDC-biotin hydrazide reaction is complete, uncoupled biotin can be removed through dialysis, preferably dialysis against distilled water. The biotinylated HA substrate (bHA) that results comprises an EDC-biotin moiety covalently bound to the HA molecule at a ratio of 1 EDC-biotin moiety for every 200 disaccharide moieties, preferably every 150 disaccharide moieties, more preferably every 85 to 100 disaccharide units in the HA molecule. The ratio of EDC-biotin to HA molecule can range from as high as 1:1 (e.g., one EDC-biotin moiety bound to a disaccharide unit) to 5:1 (e.g., one EDC-biotin moiety bound to an HA molecule containing 100 disaccharide units). The bHA reagent can be stored, preferably at -20°C , until use.

Assay devices for detection of Hase activity are prepared by coupling bHA to, for example, the surface of a microtiter well having a covalently attached moiety NH-CH_3 (e.g., Covalink-NH microtiter plates, NUNC, Placerville, NJ). An exemplary chemical and physical configuration of surface for covalent attachment is shown in Fig. 2. Methods for preparation of such substrates and the use of such substrates to covalently immobilize compounds are well known in the art (see, e.g., U.S. Patent No. 5,427,779, issued June 27, 1996, and PCT published application number WO 8905329, published June 15, 1989). Coupling

of bHA to the assay device is accomplished by, for example, incubation of bHA in the Covalink-NH well with EDC according to the manufacturer's specifications. Unbound bHA is removed by washing the wells with buffer. bHA that remains in the well of the plate is covalently bound to the surface of the well. The
5 resulting assay device comprises bHA covalently bound to the surface to be exposed to the test sample such that the bHA is bound to the surface by at least one covalent bond, preferably at least one covalent bond per every 200 disaccharide units in the bHA molecule, more preferably at least one covalent bond per 100 disaccharide units, most preferably at least one covalent bond per 50 disaccharide
10 units in the bHA molecule. The plates are preferably used within about 1 week to 10 days after preparation.

The Hase activity assay is performed by placing a test sample in the well of the assay device having covalently-bound bHA, and incubating the sample for a period sufficient to allow for hpHase in the test sample to degrade bHA in the
15 well. The test sample can be any sample suspected of having Hase activity (e.g., a biological sample (e.g., blood, serum, plasma, urine, or a sample derived from a recombinant source) or a sample from a step during biochemical purification). Preferably, a sample having a known amount of hpHase activity is subjected to the assay as a control.

20 The pH of the incubation buffer can be adjusted according to the desired pH optima of the Hase activity in the sample (e.g., neutral Hase activity (pH of about 7.0 to 7.5) or acid active Hase activity (pH below about 4.5, preferably at about pH 3.0 to 3.7)). Following incubation, the wells are washed to remove degraded bHA and remaining, undegraded bHA is detected, e.g., by reaction the remaining
25 bHA with avidin peroxidase and detection with a microplate reader. Covalent binding of bHA to the assay device prevents the undegraded bHA from being washed off the plate, thus increasing the sensitivity and accuracy of the assay. Moreover, the assay requires only 60 minutes or less to complete and is 1,000 times more sensitive than conventional colorimetric assays (Afify et al. *supra*) and
30 about 1,000 to 5,000 times more sensitive than SDS-based gel zymography. The Hase assay of the invention can be used in a variety of applications where one

desires to determine (qualitatively or quantitatively) the presence of acid active or neutral Hase activity in a sample. In one embodiment, the Hase activity assay of the invention is useful in the identification of patients having a defect in Hase activity (e.g., a LuCa-1/hpHase defect associated with reduced hpHase plasma activity). determine , and/or .

Specific activity of hyaluronidase is expressed in turbidity reducing units (TRU). One TRU is defined as the amount of hyaluronidase activity required to reduce the turbidity of an acidified solution of hyaluronan and is equivalent to the U.S.P./National Formulary (NF XIII) units (NFU). The results using the assays described above are related to the TRU, the NFU, and U.S.P. units through a standard curve of a sample of hyaluronidase (e.g., WYDASE®, Wyeth-Ayerst) standardized through the U.S.P. For example, a standard curve can be generated through co-incubation of serial dilutions of bovine testicular hyaluronidase (WYDASE®) and activity of unknown samples interpolated through a four parameter curve fit to yield values in relative TRU (rTRU)/ml (Dorfman et al., 1948, *J. Biol. Chem.* 172:367).

Biochemical hpHase Purification Method

Acid active hyaluronidase activity can be significantly enriched and/or purified using temperature-induced detergent phase extraction with a non-ionic detergent (e.g., Triton X-114). In general, a sample comprising or suspected of comprising an acid active Hase (aaHase) such as hpHase is dissolved in a solution comprising a non-ionic detergent at low temperature (e.g., substantially below room temperature, preferably less than about 15°C, more preferably about 4°C). The sample can be, for example, raw human plasma, outdated human plasma without platelet degranulation (e.g., through addition of citrate), a lipid fraction of human plasma, human blood, human serum, human urine, or conditioned medium or lysates from cells expressing recombinant aaHase (e.g., mammalian, insect, bacterial, or yeast cells, preferably mammalian cells). Preferably, the sample is human plasma or human urine, which is a particularly rich source of hpHase. Purification from plasma is advantageous over purification

from whole blood or serum since the plasma fraction contains less total protein than either serum or whole blood.

After the sample is dissolved, the temperature of the solution is raised to at least room temperature or above (preferably above about 25°C, more preferably about 37°C), thereby resulting in formation of detergent-rich and detergent-poor phases. The aaHase partitions into the detergent-rich phase. The detergent-rich phase can be further enriched for aaHase by removal of the detergent-rich phase and repetition of temperature-induced detergent phase extraction. Repeating this temperature-induced detergent phase extraction three times results in at least about 10-fold, preferably at least about 20-fold, more preferably at least about 60-fold enrichment of aaHase activity relative to aaHase activity in the starting material. The aaHase activity of the detergent-rich phase can be further enriched and purified by, for example, cation exchange chromatography and/or hydroxylapatite resin.

15

Generation and Identification of Anti-Native aaHase Antibodies

Although there are known procedures for producing antibodies from any given antigen, previous attempts to produce anti-native aaHase antibodies (e.g., anti-native hpHase antibodies) have failed. Although antibodies that bind denatured anti-aaHase have been generated, these antibodies, generated using conventional methods and a conventional ELISA assay, did not bind native (i.e., non-denatured) aaHase (Harrison et al. 1988 *J Reprod Fertil* 82:777-85). By following the procedures described herein, antibodies that bind native aaHase enzymes (e.g., native hpHase) have been obtained. Likewise, the ordinarily skilled artisan can follow the procedures outlined herein to generate other anti-native aaHase antibodies, including other anti-native hpHase antibodies. In general, the invention overcomes the problems associated with production of aaHase antibodies by providing a screening assay that detects antibodies that bind native aaHase.

Generation of anti-hpHase antibodies

aaHase can be used as an antigen in the immunization of a mammal (e.g., mouse, rat, rabbit, goat) and production of hybridoma cell lines according to methods well known and routine in the art (see, for example, Harlow and Lane, 1988, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY; Schrier et al., 1980, Hybridoma Techniques, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY). The aaHase used in the antigenic preparation can be purified from the source in which it naturally occurs (e.g., hpHase purified from human plasma), recombinant aaHase, biologically active (e.g., antigenic and/or enzymatically active) aaHase polypeptides, native aaHase, and/or denatured aaHase polypeptides. Where the aaHase is a recombinant aaHase, the recombinant aaHase can comprise the amino acid sequence of a naturally-occurring aaHase, or can be modified relative to native aaHase (e.g., by amino acid substitution, deletion, or addition (e.g., fusion protein)). Preferably, the antigenic preparation is native aaHase (e.g., aaHase purified from the source in which it naturally occurs or recombinant, full-length aaHase). aaHase can be any acid active hyaluronidase; preferably the aaHase is hpHase.

Assay for anti-aaHase antibodies

The antibodies secreted by the hybridoma cell lines are screened in the anti-native aaHase antibody assay of the invention. In general, the assay involves an insoluble support (e.g., the surface of a well of a microtiter plate) to which is bound: 1) an anti-antibody; and 2) detectably labeled hyaluronic acid (HA). The anti-antibody is capable of binding antibodies produced by the aaHase-immunized mammalian host regardless of antigen specificity. For example, where the immunized host is a mouse, the anti-antibody is a goat anti-mouse antibody (i.e., an antibody from a goat immunized with mouse antibodies). Preferably, the anti-antibody binds an Fc portion of antibodies produced by the aaHase-immunized mammalian host, and may specifically bind immunoglobulin classes or subclasses (e.g., specifically bind IgG or an IgG subclass such as IgG₁ or IgG₂). Preferably, the anti-antibody is covalently bound to the surface of the insoluble support. The

detectably labeled HA is preferably biotinylated HA (bHA) in the above-described Hase assay, and is preferably covalently bound to the surface of the insoluble plate as described above.

A schematic of the anti-aaHase assay of the invention is shown in Fig. 3.

- 5 The anti-aaHase antibody assay of the invention takes advantage of the inventors' observation that aaHases do not bind their HA substrates under non-acidic conditions (i.e., under conditions in which the aaHase is not enzymatically active). In general, the assay is performed by contacting the candidate antibody with a sample comprising native aaHase (e.g., native hpHase) to allow for formation of native aaHase/antibody complexes. Preferably, this contacting step is performed at a non-acidic, preferably neutral, pH. The sample is then contacted under non-acidic (preferably neutral) conditions with the insoluble support having bound anti-antibody and detectably labeled HA to allow for formation of native aaHase/antibody/anti-antibody complexes by binding of the anti-antibody to the candidate antibody. Preferably, excess or unbound material is washed away with a non-acidic (preferably neutral) solution.
- 10
- 15

- The wash buffer is replaced with an acidic solution having a pH that allows for enzymatic activity of the aaHase. Preferably the acidic solution has a pH that approximates the optimum pH for Hase activity of the aaHase. For example, where the aaHase is hpHase, the acidic solution preferably has a pH of 3.7. The sample is incubated with the insoluble support for a time sufficient for degradation of the detectably labeled HA by the immunoprecipitated aaHase bound in the native aaHase/antibody/anti-antibody complex, preferably about 60 min. The samples are then washed to remove degraded HA and undegraded HA is detected by virtue of its label. For example, where the detectable label is biotin, undegraded bHA is detected as described in the above-described Hase assay. Degradation of HA is correlated with the presence of aaHase in the sample which in turn is correlated with the presence of an anti-native aaHase antibody. The general characteristics of antibodies of the invention are described below.
- 20
- 25

Antibody/antigen binding forces

The forces that hold an antigen and antibody together can be classified into four general areas: (1) electrostatic; (2) hydrogen bonding; (3) hydrophobic; and (4) Van der Waals. Electrostatic forces are due to the attraction between oppositely charged ionic groups on two protein side-chains. The force of attraction (F) is inversely proportional to the square of the distance (d) between the charges. Hydrogen bonding forces are due to formation of reversible hydrogen bridges between hydrophilic groups such as -OH, -NH₂ and -COOH. These forces are largely dependent upon close positioning of two molecules carrying these groups. Hydrophobic forces operate in the same way that oil droplets in water merge to form a single large drop. Accordingly, non-polar, hydrophobic groups such as the side-chains on valine, leucine and phenylalanine tend to associate in an aqueous environment. Lastly, Van der Waals are forces created between molecules by interaction between the external electron clouds. Further information about the different types of forces is known in the art (see, e.g., *Essential Immunology*, I.M. Roitt, ed., 6th Ed. Blackwell Scientific Publications, 1988).

Useful antibodies of the present invention exhibit all of these forces. By obtaining an accumulation of these forces in greater amounts, it is possible to obtain an antibody that has a high degree of affinity or binding strength to native aaHase, and in particular an antibody that has a high degree of binding strength to aaHase in the material in which it naturally occurs (e.g., human plasma).

Measuring antibody/antigen binding strength

The binding affinity between an antibody and an antigen is an accumulative measurement of all of the forces described above. Standard procedures for carrying out such measurements are known in the art and can be directly applied to measure the affinity of anti-native aaHase antibodies of the invention.

One standard method for measuring antibody/antigen binding affinity uses a dialysis sac, composed of a material permeable to the antigen but impermeable to the antibody. Antigens that bind completely or partially to antibodies are placed within the dialysis sac in a solvent (e.g., water). The sac is then placed within a larger container which does not contain antibodies or antigen but contains only the

solvent. Since only the antigen can diffuse through the dialysis membrane the concentration of the antigen within the dialysis sac and the concentration of the antigen within the outer larger container will attempt to reach an equilibrium. The amount of antigen that remains bound to antibody in the dialysis sac and the amount that disassociated from the antibody are calculated by determining the antigen concentrations within the dialysis sac and within the solvent outside the dialysis sac. By constantly renewing the solvent (e.g., the water) within the surrounding container so as to remove any diffused antigen, it is possible to totally disassociate the antibody from antigen within the dialysis sac. If the surrounding solvent is not renewed, the system will reach an equilibrium, and the equilibrium constant (K) of the reaction, i.e., the association and disassociation between the antibody and antigen, can be calculated. The equilibrium constant (K) is calculated as an amount equal to the concentration of antibody bound to antigen within the dialysis sac divided by the concentration of free antibody combining sites times the concentration of free antigen. The equilibrium constant or "K" value is generally measured in terms of liters per mole. The K value is a measure of the difference in free energy (ΔG) between the antigen and antibody in the free state as compared with the complexed form of the antigen and antibody. Anti-native aaHase antibodies having an affinity or K value of 10^7 l/mole to 10^9 l/mole or more are preferred.

Antibody avidity

As indicated above the term "affinity" describes the binding of an antibody to a single antigen determinate. The term "avidity" is used to express the interaction of an antibody with a multivalent antigen. The factors that contribute to avidity are complex and include both the heterogeneity of the antibodies in a given serum that are directed against each determinate on the antigen and the heterogeneity of the determinants themselves. The multivalence of most antigens leads to an interesting "bonus" effect in which the binding of two antigen molecules by an antibody is always greater, usually many fold greater, than the arithmetic sum of the individual antibody links. Thus, it can be understood that the measured avidity between an antiserum and a multivalent antigen will be

somewhat greater than the affinity between an antibody and a single antigen determinate.

Uses of Anti-Native aaHase Antibodies

5 Anti-native aaHase antibodies are useful in various immunotechniques, including immunopurification and immunodetection techniques. Anti-native aaHase antibodies useful in such immunotechniques may be either polyclonal or monoclonal antibodies, preferably monoclonal antibodies.

10 Preferably, anti-native aaHase antibodies useful in immunotechniques exhibit an equilibrium or affinity constant (K_d) of at least 10^7 l/mole to 10^9 l/mole or greater. The binding affinity of 10^7 l/mole or more may be due to (1) a single monoclonal antibody (i.e., large numbers of one kind of antibody) (2) a plurality of different monoclonal antibodies (e.g., large numbers of each of five different monoclonal antibodies) or (3) large numbers of polyclonal antibodies. It is also
15 possible to use combinations or (1)-(3).

20 Preferred antibodies bind 50% or more of native aaHase in a sample. However, this may be accomplished by using several different antibodies as per (1)-(3) above. An increased number of different antibodies is generally more effective than a single antibody in binding a larger percentage of antigen in a sample. Thus, a synergistic effect can be obtained by combining combinations of two or more antibodies which bind native aaHase, i.e., by combining two or more
20 antibodies that have a binding affinity K_a for native aaHase of 10^7 l/mole or more.

Immunopurification using anti-native aaHase antibodies

25 Anti-native aaHase antibodies can be used in, for example, immunopurification of aaHase from its naturally occurring source (e.g., hpHase from human blood, plasma, serum, or urine) or from a source of recombinant aaHase production (e.g., from supernatants or cell lysates of transformed cells expressing hpHase). Immunopurification techniques useful with the anti-native
30 aaHase antibodies of the invention include, but are not limited to, immunoprecipitation, immunoaffinity isolation on beads, immunoaffinity column

chromatography, and other methods well known in the art. Anti-native aaHase antibodies useful in immunopurification techniques. The immunopurification methods using the antibodies of the invention can use a single anti-native aaHase antibody (e.g., a monoclonal or polyclonal antibody, preferably a monoclonal antibody) or multiple anti-native aaHase antibodies.

In addition, the anti-native aaHase antibodies of the invention can be used to prepare a device for immunopurification of native aaHase, preferably native hpHase. In general, such devices are prepared by covalently binding an anti-native aaHase antibody to an insoluble support (e.g., bead, affinity column component (e.g., resin), or other insoluble support used in immunoaffinity purification). Alternatively, the antibody may be bound to a metal particle which allows separation of anti-native aaHase-aaHase complexes from a solution by use of a magnetized column. The anti-aaHase antibody may be a monoclonal or polyclonal antibody, preferably a monoclonal antibody. The antibodies bound to the insoluble support (or otherwise employed in purification) may also comprise a mixture of anti-native aaHase antibodies to provide a device that can bind to at least 50% of the native aaHase in the sample. Such immunopurification devices can be used to isolate aaHase from a source in which it naturally occurs (e.g., hpHase from raw serum, raw plasma, or urine) or from a source of recombinantly-produced aaHase.

Qualitative and quantitative immunodetection using anti-native aaHase antibodies

Anti-native aaHase antibodies can be used in immunodetection assay to detect and, where desirable, quantitate aaHase in a sample. Immunodetection assays using anti-native aaHase antibodies can be designed in a variety of ways. For example, anti-native aaHase antibodies can be used to produce an assay device comprising anti-native aaHase antibodies bound to a soluble support (e.g., an immunoassay column, beads, or wells of a microtiter plate). Methods for covalent or non-covalent attachment of an antibody to a soluble support are well known in the art. A sample suspected of containing an aaHase is then contacted with the assay device to allow formation of anti-native aaHase antibody-aaHase complexes.

The anti-native aaHase-aaHase complexes can then be detected by virtue of an aaHase activity associated with the complex as described in the anti-native aaHase assay described above, or by contacting the complex with a second detectably-labeled anti-native aaHase antibody.

5 By "detectably labeled antibody", "detectably labeled anti-aaHase" or "detectably labeled anti-aaHase fragment" is meant an antibody (or antibody fragment that retains antigen binding specificity), having an attached detectable label. The detectable label is normally attached by chemical conjugation; where the label is a polypeptide, the label can be attached by genetic engineering
10 techniques. Detectable labels may be selected from a variety of such labels known in the art, but normally are radioisotopes, fluorophores, paramagnetic labels, enzymes (e.g., horseradish peroxidase), or other moieties or compounds that either emit a detectable signal (e.g., radioactivity, fluorescence, color) or emit a detectable signal after exposure of the label to its substrate. Various detectable
15 label/substrate pairs (e.g., horseradish peroxidase/diaminobenzidine, avidin/streptavidin, luciferase/luciferin), methods for labelling antibodies, and methods for using labeled antibodies are well known in the art (see, for example, Harlow and Lane, eds. (*Antibodies: A Laboratory Manual* (1988) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY)).

20 Alternatively, detectably labeled anti-native aaHase antibodies can be directly used to detect and/or quantify aaHase in a sample. For example, detectably labeled anti-native aaHase antibodies can be contacted with a tissue sample suspected of having a LuCa-1/hpHase defect (e.g., a tissue sample derived from breast, ovaries, or lung) for a time sufficient to allow for formation of
25 complexes between the anti-native hpHase antibody and hpHase in the tissue sample (e.g., hpHase in the plasma membrane of cells of the tissue sample). Binding of the anti-native hpHase antibody can then be detected and/or quantified by virtue of a detectable label bound to the anti-native hpHase antibody. Alternatively, binding of the anti-native hpHase antibody can be detected using an
30 antibody that binds the anti-native hpHase antibody. Binding of anti-native hpHase antibody to a tissue sample can then be compared to anti-native hpHase

antibody binding to a control sample (e.g., a normal sample having no LuCa-1/hpHase defect and/or a sample containing tissue associated with a LuCa-1/hpHase defect) and the antibody binding correlated with the presence or absence of a LuCa-1/hpHase defect in the patient.

- 5 The aaHase immunodetection assays can be used in a variety of different ways with a variety of samples, as will be apparent to one of ordinary skill in the art upon reading the disclosure provided herein. For example, the aaHase immunodetection assays can be used to detect hpHase in serum or plasma samples of patient receiving hpHase therapy and/or to correlate hpHase in the serum or
10 plasma of a patient with tumor progression in the patient, responsiveness to therapy, and/or uptake of hpHase by the patient's body.

Where hpHase is detected in serum or urine, the levels of hpHase can be correlated with susceptibility to and/or the presence of a LuCa-1 defect associated disease state and/or the severity of such disease. For example, hpHase levels in
15 patients having lung cancer associated with a heterozygous defect in LuCa-1 are approximately 50% of normal hpHase levels, while patients having a homozygous defect in LuCa-1 exhibit hpHase levels that are very low or undetectable. Thus, hpHase levels, as detected in an aaHase assay of the invention or detected using an anti-native hpHase antibody of the invention, can not only be correlated with
20 tumor progression, but can also be directly correlated with a genetic defect in LuCa-1, thus allowing identification of patients susceptible to conditions associated with LuCa-1 defects, e.g., cancer. Thus, the present invention allows one to directly correlate hpHase levels with the number of functional alleles in a patient and to detect a genetic defect by simply ascertaining the level of hpHase in blood,
25 plasma, serum or urine.

Methods of Making hpHase

In addition to the purification procedure outlined above, hpHase hyaluronidase polypeptides can be made by standard synthetic techniques, or by
30 using recombinant DNA technology and expressed in bacterial, yeast, or mammalian cells using standard techniques. As used herein, the term "hpHase"

includes natural, recombinant, and modified forms of the protein unless the context in which the term is used clearly indicates otherwise.

Chemical Synthesis

hpHase polypeptides can be synthesized based on the amino acid sequences described herein and variations thereof by standard solid-phase methods using the *tert*-butyloxy-carbonyl and benzyl protection strategy described in Clark-Lewis et al., *P.N.A.S., USA*, 90:3574-3577 (1993) and Clark-Lewis et al., *Biochemistry*, 30:3128-3135 (1991). After deprotection with hydrogen fluoride, the proteins are folded by air oxidation and purified by reverse-phase HPLC. Purity is determined by reverse-phase HPLC and isoelectric focusing. Amino acid incorporation is monitored during synthesis, and the final composition is determined by amino acid analysis. The correct covalent structure of the protein can be confirmed using ion-spray mass spectrometry (SCIEX APIII).

Recombinant DNA Techniques for Synthesis of hpHase Polypeptides

As discussed in the examples below, LuCa-1 (SEQ ID NO:3) and hpHase (SEQ ID NO:1) are identical. The only variation between the amino acid sequence of hpHase (SEQ ID NO:1) and the amino acid sequence of LuCa-1 (SEQ ID NO:3) is a substitution of Val for Leu in the N-terminus at the 27th amino acid residue (where the Met at the N-terminus of the signal sequence is counted as the first amino acid residue); however, the two proteins are otherwise identical in amino acid sequence and immunological and biochemical characteristics. The nucleotide sequence encoding hpHase is identical to the sequence encoding LuCa-1 (SEQ ID NO:4), except that the cytosine of the third residue in the codon corresponding to the 27th amino acid residue (Leu in LuCa-1; Val in HpHase) is substituted with guanine. Thus, the nucleotide sequence encoding LuCa-1 (SEQ ID NO:4) is the nucleotide sequence encoding hpHase. The LuCa-1/hpHase gene has been isolated and sequenced (Bader et al. GenBank accession no. U03056, NID G532973, submitted Nov. 1, 1993; see also GenBank Accession No. U96078). The amino acid and nucleotide sequences of LuCa-1 as described by Bader et al. are provided below.

MAGHLLPICALFLTLLDMAQGFGRGPLVNPFPFTTVWNANTQWCLERHGVDVDVSVFVDV
 VANPGQTFRGPDMTIFYSSQLGTYPPYTPTGEPVFGGLPQNASLIAHLARTFQDILAAIPAP
 DFSGLAVIDWEAWRPRWAFNWDTKDIYRQRSRALVQAQHPDWPAPQVEAVAQDQFQGA
 ARAWMAGTLQLGGALRPRGLWGFYGFDPDCYNYDFLSPNYTGQCPSGIRAQNDQLGWLW
 5 GQSRALYPSIYMPAVLEGTGKSQMYVQHRVAEAFRVAVAAGDPNLPVLPYVQIFYDTTN
 HFLPLDELEHSLGESAAQGAAGVVLWVSWENTRTKESCQAIKEYMDTTLGPFILNVTSGA
 LLCSQALCSGHGRCVVRTSHPKALLLNPAFSIQLTPGGGPLSLRGALSLEDQAQMAVEF
 KCRCPGWQAPWCERKSMW (SEQ ID NO:3)

10 1 ttctccagg agtctctggt gcagctgggg tggaaatctgg ccaggccctg cttaggcccc
 61 catcttgggg tcaggaaatt tggaggataa ggcccttcag cccaagggtc agcaggggacg
 121 agcggggcaga ctggcgggtg tacaggaggg ctgggtgac ctgtcttgg tcaactaggc
 181 catggatct tctccagg gctgccagga ttctgggtg aagagacagg aaggctccc
 241 ccccttggc gggtcagct gggggctgag ggctggctg tcagccactc ttccagaac
 15 301 atatgcatg gcctcagtg ctcaggggga agcagggggtg ggcgagctta ggclagagca
 361 agtctctgg gagatggcag aggcctggc tgaggggcaa ctggatgtg cctccagtg
 421 gccatgtcc cctccatgc tctccctgc cctctggag cctgcaggt caatgttaa
 481 cagaaccag agcagcgtg gattaatgc caagggtca gccccagc cctgagcagt
 541 gggggaatg gagacttgc aacctgtct cagctctgc tccctgggc aggtgtctt
 20 601 cgaccagtc cgtgccatg caggccacct gctccctc tggccctc tctgacct
 661 actcgatag gcccaaggct ttagggggcc ctggtaacc aaccggcct tcaccaccgt
 721 ctggaatga aacaccagt ggtgctgga gaggcacgt gggacgtg atgtcagt
 781 ctctgatgt gtagcaacc cagggcagac ctccgggc cctgacatg caatttcta
 841 tagtccag ctggcacct accctacta cagccactc ggggagcctg tgttgggtg
 25 901 tctgcccag aatgccagc tgattccca cctggcccgc acattccag acatctggc
 961 tggcatact gctctgact tctcagggt ggcagtcac gactggagg catggccc
 1021 acgtgggc ttaactggg acaccaagga cattaccg cagcctcag ggcactgt
 1081 acaggagcag caccctgatt ggccagctcc tcaggtggag gcagtagccc aggaccagt
 1141 ccaggagct gcacgggct ggatggcagg caccctccag ctggggggg cactgcgtc
 30 1201 tggggctct tggggtct atggtctcc tgaactac aactatgact tttaagccc
 1261 caactacac gccagtgcc catcaggcat ccgtgccc aaatgaccagc taggtggct
 1321 gtggggccag agccgtgccc tctatccag catctacag cccgagctg tggaggccac
 1381 agggaaatga cagatgatg tgcaacaccg tgggcccag gcatccgtg tggctgtggc
 1441 tgcgtgtgac ccaatctgc cgtgtctgc ctatgtccag atctctatg acacgacaaa
 35 1501 ccactttct cccctggatg agctggagca cagcctggg gagagtgcg cccagggggc
 1561 agctggagt gtgtctggg tgaactggga aaatacaga accaaggaa catgtcaggc
 1621 catcaaggag tatatggaca ctacactgg gcccttcac ctgaactga ccagtgggc
 1681 ccttctctgc agtcaagccc tgtgtccgg ccagggcgc tgtgtccgc gcaccagcca
 1741 ccccaagcc cctctctcc ttaacctgc cagtctcc atccagctca cgcctgttg
 40 1801 tggccccctg agcctgcggg gtgccctc actgaagat caggcacaga tggctgtga
 1861 gttaaatgt cgtatgacc ctggctggca ggcaccgtg tgtagcggga agagcatgt
 1921 gtgattggc acacactgag ttgcacat tgagaacct atgcactct ggtctggcca
 1981 gggcttctc aaatacatg acagtcatac aagtcaggt cacagtaaag agtacactca
 2041 gccacttca caggcatat cctgcacac acatgcatac ttacagactg gaatagtggc
 45 2101 ataaggagt agaaccacag cagacaccat tcatctgc tcatatgca tctacttgc
 2161 aaggctatg acaattctc cagagacact gaggcagct tgaactgca gcaatcaca
 2221 aggtgacat tcatgagt cctactctt gccaatccc gtgtaagcg ttatgttg
 2281 acttaattc tctcacaat gaggctatga ggaactgag tcatcaca tgagagtaag
 2341 cagctggcc aggtgtcac agcaaaaaa gggagaagt gagattcaa cccagctgt
 50 2401 ctagctccg ggtacagcc ctgcactc tactgagtt gtgtaacca gccctgcag
 2461 accctgaat ctgtgagag gcaccagtc agcaataaa gcagtcata ttactt (SEQ ID NO:4)

The nucleotide sequence encoding hpHase can be isolated according to any one of a variety of methods well known to those of ordinary skill in the art. For example, DNA encoding hpHase can be isolated from either a cDNA library or from a genomic DNA library by hybridization methods. Alternatively, the DNA
5 can be isolated using standard polymerase chain reaction (PCR) amplification of synthetic oligonucleotide primers, e.g., as described in Mullis et al., U.S. Patent No. 4,800,159, or expression cloning methods well known in the art (see, e.g., Sambrook et al. Sambrook et al. 1989 Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY). Where
10 hybridization or PCR is used to identify DNA encoding hpHase, the sequence of the oligonucleotide probes or primers can be based upon the amino acid or nucleotide sequence of LuCa-1 provided above. The sequence of isolated hpHase polypeptide-encoding DNA can be determined using methods well known in the art (see, for example, Sambrook et al., supra). Following sequence confirmation, the
15 resulting clones can be used to for example, identify homologs of a hpHase (e.g., other human alleles encoding hpHase or an acid active serum hyaluronidase of another mammalian species (e.g., dog, rat, mouse, primate, cow), and/or to transform a target cell in a host for expression of the hpHase-encoding DNA (e.g., as in anti-cancer therapy through hpHase polypeptide expression in host cells,
20 preferably cancerous cells of a patient).

Production of hpHase-Encoding Constructs and Expression of hpHase Polypeptides

Numerous vectors are available for production of hpHase-encoding
25 constructs and hpHase expression (see, e.g., the American Type Culture Collection, Rockville, MD). Preferably the vector is capable of replication in both eukaryotic and prokaryotic hosts, and are generally composed of a bacterial origin of replication and a eukaryotic promoter operably linked to a DNA of interest, thereby allowing production of hpHase-encoding constructs. Suitable host cells,
30 as well as methods for constructing stably-transformed host cell lines, are also publicly available, e.g., Pouwels et al., 1985, *Cloning Vectors: A Laboratory*

Manual, Ausubel et al., 1989, *Current Protocols in Molecular Biology*, John Wiley & Sons, New York; Sambrook et al., *supra*; Kormal et al., 1987, *Proc. Natl. Acad. Sci. USA*, 84:2150-2154; each of which are hereby incorporated by reference with respect to methods and compositions for manipulation of a DNA of interest).

5

Expression of an hpHase polypeptide is accomplished by inserting a nucleotide sequence encoding an hpHase polypeptide into a nucleic acid vector such that a promoter in the construct is operably linked to the hpHase-encoding sequence, which construct is then used to transform a host cell. hpHase
10 expression can be accomplished in a mammalian cell line by either transient, constitutive, or inducible expression. In one preferred embodiment, the host cell is a mammalian cell, preferably a cos-7 cell line, or a cos-7 cell-derived cell line. More preferably, the host cell for hpHase production is a mammalian cell that is protected against hpHase expression (i.e., the cell can produce high levels of
15 hpHase without adverse affects upon the host cell (e.g., slow cell growth, cell death)). In one embodiment, the hpHase-resistant cell is an SV-transformed cell line or an adenovirus-transformed cell line (i.e., a cell line transformed with sheared adenovirus), preferably the HEK cell line (HEK 293). For example, transformation of HEK cells with hpHase-encoding DNA driven by a strong
20 promoter (e.g., CMV) provides surprisingly high hpHase expression and secretion into the culture medium (approximately 2.9×10^{-10} mg/cell/24 hrs.) (> 100 rTRU). Use of such cell lines allows for production of sufficient quantities of hpHase for use in, e.g., protein therapy.

Where expression is desired in a tumor cell line that is not resistant to
25 hpHase, the hpHase-encoding DNA is preferably under control of an inducible promoter, preferably an inducible promoter responsive to an inducing agent that does not significantly affect other mammalian cells in the vicinity of the transformed, hpHase construct-containing cells. For example, the hpHase-encoding DNA can be under the control of a steroid inducible promoter (e.g., in
30 the ecdysone expression system, which is inducible by muristerone). The target cells are transformed and, upon exposure to the inducing agent, express hpHase.

Where the hpHase-encoding DNA is used to transform tumor cells for treatment of a tumor-bearing patient, the hpHase-encoding DNA may be transiently, constitutively, or inducibly expressed, preferably transiently or
5 constitutively expressed, more preferably constitutively expressed. In addition or alternatively, peritumor cells (e.g., cells adjacent the tumor cells) or other cells capable of expressing and secreting hpHase (e.g., liver cell or monocyte) can be transformed.

Recombinant hpHase polypeptide expression (e.g., produced by any of the
10 expression systems described herein) can be assayed by immunological procedures, such as Western blot or immunoprecipitation analysis of recombinant cell extracts, or by the Hase activity assay of the invention as described herein. For example, hpHase polypeptides according to the invention can be produced by transformation of a suitable host cell with an hpHase polypeptide-encoding nucleotide sequence
15 in a suitable expression vehicle, and culturing the transformed cells under conditions that promote expression of the encoded polypeptide, and preferably secretion of hpHase into the culture medium. The method of transformation and the choice of expression vehicle will depend on the host system selected. Those skilled in the field of molecular biology will understand that any of a wide variety
20 of prokaryotic and eukaryotic expression systems may be used to produce hpHase polypeptides of the invention.

Identification of Biologically Active hpHase Polypeptides

hpHase polypeptide-encoding DNAs can encode all or a portion of an
25 hpHase. Preferably, the hpHase polypeptide is biologically active, e.g., exhibits acid active hyaluronidase activity in the cleavage of hyaluronan and/or can be bound by an anti-native hpHase antibody. In general, once information regarding the ability of a protein to elicit antibodies and/or information regarding an enzymatic or other biological activity of a protein of interest is known, methods
30 for identification of biologically active polypeptides of the full-length protein are routine to the ordinarily skilled artisan, particularly where the nucleotide sequence

and/or amino acid sequence encoding the protein of interest (here hpHase) is provided as in the present case.

Biologically active hpHase polypeptides can be identified by using the Hase activity assay of the invention, or by using conventional Hase activity assays (e.g., the ELISA-like hyaluronan assay (Stern et al., 1992, *Matrix* 12:391-403) or substrate gel zymography (Guentenhoener et al., 1992, *Matrix* 12:388-396)). Alternatively, biologically active hpHase polypeptides can be detected by binding of an anti-native hpHase antibody to a component of the transformed host cell supernatant and/or lysate. hpHase polypeptides preferably exhibit at least 25%, more preferably 50%, still more preferably 75%, even more preferably 95% of the activity of native hpHase.

Identification of Hyaluronidases Homologous to hpHase

DNA encoding hyaluronidases homologous to hpHase (e.g., contain conservative amino acid substitutions relative to a native hpHase) can be accomplished by screening various cDNA or genomic DNA libraries by hybridization or PCR using oligonucleotides based upon the DNA sequence and/or amino acid sequence of an hpHase (e.g., a LuCa-1/hpHase sequence). Alternatively the oligonucleotides used may be degenerate, e.g., based upon a selected amino acid sequence of hpHase or designed so as to allow detection or amplification of DNA encoding an hpHase-like amino acid sequence having conservative amino acid substitutions and/or to take into account the frequency of codon usage in the mammalian species DNA to be screened. Such "degenerate oligonucleotide probes" can be used in combination in order to increase the sensitivity of the hybridization screen, and to identify and isolate hpHase analogs in other species or variant alleles encoding hpHase in humans. Methods for designing and using degenerate oligonucleotide probes to identify a protein for which an amino acid and/or nucleotide sequence, as well as methods for hybridization and PCR techniques for screening and isolation of homologous DNAs, are routine and well known in the art (see, for example, Sambrook et al. *supra*).

Alternatively, the DNA encoding hpHase may be isolated by expression cloning methods well known in the art (see, for example, Sambrook et al., supra). For example, mammalian cells can be transformed with a cDNA expression library, i.e., a collection of clones containing various cDNA fragments operably
5 linked to a eukaryotic promoter. Expression of an hpHase homology or a biologically active fragment thereof can be detected by assaying the culture supernatant and/or cell lysates using the Hase activity assay of the invention.

Therapies Using hpHase Polypeptides

10 The substantially pure native hpHase polypeptides (e.g., hpHase polypeptides that are not associated with the components of plasma from which hpHase is purified) of the invention can be used in a variety of applications including human and veterinary therapies, either alone or in combination with other therapeutic agents. Purified hpHase of the invention can generally be used in
15 place of neutral Hase formulations or WYDASE™, where the condition to be treated is associated with excess hyaluronic acid and/or therapy is designed to increase Hase activity generally (i.e., the conventional neutral hyaluronidase-containing formulation is not used to treat a specific defect in neutral Hase activity, but rather provides a Hase (neutral or acid active) activity). Use of acid
20 active hpHase is preferred to use of neutral Hases since acid active hpHase can provide controlled degradation of HA substrate and does not degrade all components of the extracellular matrix in the patient.

hpHase can be used in the treatment of diseases associated with excess hyaluron, to enhance circulation of physiological fluids at the site of administration
25 (e.g., as a spreading agent, e.g., by subcutaneous or topical application (e.g., in cosmetic formulations such as cosmetic creams), and/or as an anti-cancer agent either alone or in combination with chemotherapeutic agents. For example, hpHase can be administered to a patient to facilitate clysis, particularly hypodermoclysis. hpHase can also be administered to patients suffering from
30 stroke or a myocardial infarction (e.g., by infusion). Preferably, hpHase is administered in the absence, or at very low levels, of heparin, a powerful inhibitor

- of hyaluronidase. Methods for administration, and amounts of hpHase administered, for treatment of myocardial infarction can be based upon methods of administration of bovine testicular hyaluronidase and amounts administered (see, e.g., Wolf et al. 1982 *J. Pharmacol. Exper. Therap.* 222:331-7; Braunwald et al. 5 1976 *Am. J. Cardiol.* 37:550-6; DeGiovanni et al. 1961 *Br. Heart J.* 45:350; DeOliveira et al. 1959 *Am. Heart J.* 57:712-22; Kloner et al. 1978 *Circulation* 58:220-6; Kloner et al. 1977 *Am. J. Cardiol.* 40:43-9; Koven et al. 1975 *J. Trauma* 15:992-8; Maclean et al. 1978 *J. Clin. Invest.* 61:541-51; Maclean et al. 1976 *Science* 194:199-200; Maroko et al. 1975 *Ann. Intern. Med.* 82:516-20; 10 Maroko et al. 1977 *N. Engl. J. Med.* 296:896-903; Maroko et al. 1972 *Circulation* 46:430-7; Salet 1980 *Clin. Biochem.* 13:92-94; Snell et al. 1971 *J. Clin. Invest.* 50:2614-25; Wolf et al. 1981 *Circ. Res.* 48:88-95).

Furthermore, hpHase can also have therapeutic effects when administered to patients having certain lysosomal storage diseases associated with a defect in 15 hyaluronidase (see, e.g., Natowicz et al. 1996 *N. Engl. J. Med.* 335:1029-33). hpHase can be used therapeutically by direct administration of hyaluronidase (e.g., intracellularly or intravenously) as a form of shunt pathway and/or by gene therapy (e.g., to replace defective copy(ies) of the *LuCa-1* gene). Lysosomal storage disease amenable to hpHase therapy are those diseases that result in accumulation 20 of [GlcNAc β 1-4GlcUA β 1-3]_n (GAGs) due to a defective mannose-6-phosphate pathway. hpHase can degrade these accumulated GAGs under a nonfunctional cellular system since Hase activity does not depend upon the mannose-6-phosphate pathway (Herd et al. 1976 *Proc. Soc. Experim. Biol. Med.* 151:642-9).

hpHase can also be used in the treatment of edema associated with brain 25 tumors, particularly that associated with glioblastoma multiform. The edema associated with brain tumors results from the accumulation of hyaluronan in the non-cancerous portions of the brain adjacent the tumor. Administration of hyaluronidase to the sites of hyaluronan accumulation (e.g., by intravenous injection or via a shunt) can relieve the edema associated with such malignancies 30 by degrading the excess hyaluronan at these sites. Thus, hyaluronidase is successful in the treatment of brain tumors not only in the reduction of the tumor

mass and inhibition of tumor growth and/or metastasis, but it also is useful in relieving edema associated with the malignancy. hpHase can be administered for treatment of edema in a manner similar to that for administration of bovine testicular hyaluronidase to treat edema (see, e.g., *SaEarp Arq. Braz. Med.* 44:217-20).

Of particular interest is the use of hpHase polypeptides in the treatment of metastatic and non-metastatic cancers, particularly metastatic cancers, having decreased to undetectable hpHase activity relative to non-cancerous (normal) cells. hpHase can be used as a chemotherapeutic agent (alone or in combination with other chemotherapeutics) in the treatment of any of a variety of cancers, particularly invasive tumors. For example, hpHase polypeptides can be used in the treatment of small lung cell carcinoma, squamous lung cell carcinoma, as well as cancers of the breast, ovaries, head and neck, or any other cancer associated with depressed levels of hpHase or with a defective LuCa-1 (hpHase) gene (e.g., a LuCa-1 gene that does not provide for expression of adequate hpHase levels or encodes a defective hpHase that does not provide for an adequate level of hyaluronidase activity) or other defect associated with decreased hpHase activity.

hpHase can also be used to increase the sensitivity of tumors that are resistant to conventional chemotherapy. In one embodiment, hpHase is administered to a patient having a tumor associated with a LuCa-1 defect in an amount effective to increase diffusion around the tumor site (e.g., to increase circulation of chemotherapeutic factors (e.g., to facilitate circulation and/or concentrations of chemotherapeutic agents in and around the tumor site), inhibit tumor cell motility (e.g., by HA degradation) and/or to lower the tumor cell(s) threshold of apoptosis (i.e., bring the tumor cell(s) to a state of anoikis), a state that renders the tumor cell(s) more susceptible to the action of chemotherapeutic agents or other agents that may facilitate cell death, preferably preferentially facilitate programmed cell death of cells in anoikis. Chemotherapeutic agents as used herein is meant to encompass all molecules, synthetic (e.g., cisplatin) as well as naturally-occurring (e.g., tumor necrosis factor (TNF)), that facilitate inhibition

of tumor cell growth, and preferably facilitate, more preferably preferentially facilitate tumor cell death.

Patients having or susceptible to a disease or condition that is amenable to treatment with hpHase can be identified using a variety of conventional methods, or by using an assay device of the invention having bound anti-native hpHase antibodies as described above to determine blood, plasma, serum, or urine hpHase levels, preferably blood, plasma, or serum, and correlate such levels with a LuCa-1 defect. For example, where the patient is suspected of having or of being susceptible to a condition associated with decreased hpHase activity, a biological sample (e.g., blood, serum, or plasma) can be obtained from the patient and assayed using the Hase assay and/or immunoassays using anti-native hpHase antibodies as described above.

Alternatively, particularly where the patient has or is suspected of having a cancer associated with a LuCa-1/hpHase defect, anti-native hpHase antibodies can be used to determine the levels of hpHase in a tissue sample (e.g., hpHase present in the plasma membrane of, inside of, or surrounding cells associated with the tumor tissue). Decreased levels of hpHase in the tissue sample relative to levels of hpHase associated with a normal (i.e., non-cancerous) tissue sample is indicative of a LuCa-1/hpHase defect-associated cancer.

The route of administration and amount of hpHase administered will vary widely according to the disease to be treated, and various patient variables including size, weight, age, disease severity, and responsiveness to therapy. Methods for determining the appropriate route of administration and dosage are generally determined on a case-by-case basis by the attending physician. Such determinations are routine to one of ordinary skill in the art (see, for example, Harrison's Principles of Internal Medicine, 11th Ed., 1987). For example, where hpHase is used to facilitate hypodermoclysis, a solution containing hpHase is administered by subcutaneous injection to facilitate absorption of the solution (e.g., nutrient, body fluid replacement, or blood pressure increasing solution). Preferably, hpHase is administered by injection, e.g., parenteral injection including subcutaneous, intramuscular, intraorbital, intracapsular, peritumoral, and

intravenous injection.

In a preferred embodiment, hpHase polypeptide is delivered to the patient having an hpHase-treatable tumor by delivery of hpHase polypeptide directly to the tumor site, e.g., by peritumoral injection of hpHase, injection or hpHase
5 directly into the tumor mass, and/or introduction of hpHase-encoding DNA in a tumor cell, peritumor cell, or other cell capable of expressing and secreting hpHase (e.g., liver cell or monocyte). Because hpHase has a fatty acid modification (e.g., a lipid moiety), hpHase may be readily incorporated into the plasma membrane of cells following injection at the site where hpHase action is
10 desired (see, e.g., 1995 *J. Cell Bio.* 131(3):669-77). Further, hpHase may also complex with LDL (Low Density Lipoprotein) receptors and be internalized into the cell by receptor-mediated endocytosis.

In one preferred embodiment, hpHase is delivered to the patient within a liposome formulation to provide intracellular delivery of hpHase to the target cell
15 (e.g., a cancerous cell having a LuCa-1/hpHase defect) and/or incorporating hpHase into the plasma membrane of such target cell. Preferably, the liposome formulation is prepared to provide for specific, targeted delivery of hpHase incorporated within the liposome to a target cell. Because hpHase is fairly insoluble in aqueous solution (e.g., hpHase precipitates in aqueous solution when
20 the NaCl concentration is less than 50 mM), liposome incorporation creates a more soluble preparation due to the biochemical characteristics of hpHase associated with its post-translational fatty acid chain modification. Furthermore, introduction of hpHase into the circulation would likely result in association of hpHase with high density lipoprotein (HDL) complexes, since hpHase co-purifies with these
25 lipid fractions of plasma.

The biochemical characteristics identified by the present inventors indicate that hpHase can be easily incorporated into liposomes. Methods for preparation of liposomes and administration of same are well known in the art. For example, hpHase-containing liposomes can be prepared by combining detergent-free
30 immunoaffinity purified (IAP) hpHase and liposome formulation as described in Liposome Technology, G. Gregoriadis, ed., 1984, CRC Press, Boca Raton,

Florida. Preferably the liposome formulation of the invention comprises a compound(s) that enhance hpHase activity. In a preferred embodiment, the hpHase liposome formulation comprises cholesterol and/or cardiolipin, more preferably, cardiolipin. Without being held to theory, cardiolipin enhances
5 hpHase activity by preventing the denaturation of hpHase, thus maintaining the stability of the enzyme and enhancing its incorporation into the plasma membranes of target cells. As discovered by the inventors, hpHase activity is not enhanced by the presence of phosphatidyl-ethanolamine or phosphatidyl-choline. Liposome formulations comprising cholesterol and cardiolipin are especially preferred.

10 Especially where hpHase is to be used in therapy, e.g., chemotherapy, it may be desirable to modify hpHase to provide one or more desirable characteristics. For example, although hpHase is a serum protein and thus should inherently have a substantial half-life in serum, it may be desirable to increase its biological half-life (e.g., serum half-life) by, for example, modification of the
15 polypeptide. Various methods for increasing the half-life of a protein are well known in the art and include, for example, conjugation of the protein to polyethylene glycol moieties, i.e., PEGylation (see, for example, USPN 4,179,337; USPN 5,166,322; USPN 5,206,344; Nucci et al., 1991, *Adv. Drug Delivery Rev.* 4:133-151; Zalipsky et al., 1991, "Polymeric Drugs and Drug Delivery Systems,"
20 ACS) conjugation of the protein to dextran (Maksimenko, 1986, *Bull. Exp. Biol. Med. (Russian)* 52:567-569), and deglycosylation of the protein by treatment with endoglycosidase F (Lace et al., 1990, *Carbohydrate Res.* 208:306-311).

In general, these methods are designed to increase the molecular weight of the protein, decrease the sensitivity of the protein to proteinases, and/or decrease
25 the rate of clearance of the protein from the subject to be treated. The increased half-life of modified hpHase polypeptides decreases the amount of protein needed for an effective dosage, reduces the number and frequency of administrations required, and decreases the patient's exposure to the protein, thus decreasing the potential for allergic reactions, toxic effects, or other side effect. These
30 characteristics of modified hpHase polypeptides having an increased half-life also allow for long-term use of the protein with less potential for undesirable side

effects related to protein immunogenicity and/or toxicity. Preferably these methods can be used so that the half-life of the protein is increased without substantially compromising the protein's biological activity. The enzymatic activity of hpHase can be assayed by, for example, using the Hase activity assay of the invention.

5 Methods for testing the biological half-life of a proteins are well known in the art.

The specific dosage appropriate for administration can be readily determined by one of ordinary skill in the art according to the factors discussed above (see, for example, Harrison's Principles of Internal Medicine, 11th Ed.,

10 1987). In addition, the estimates for appropriate dosages in humans may be extrapolated from determinations of the level of enzymatic activity of hpHase *in vitro* and/or dosages effective in animal studies. For example, 70-300 TRU hyaluronidase is effective in reducing the tumor load in a scid mouse. Given this information, the corresponding dosages in the average 70 kg human would range

15 from about 250,000-1,200,000 TRU hyaluronidase. The amount of hpHase polypeptide administered to a human patient is generally in the range of 1 TRU to 5,000,000 TRU of enzymatic activity, preferably between about 1,000 TRU to 2,500,000 TRU, more preferably between about 100,000 TRU to 1,500,000 TRU, normally between about 250,000 TRU and 1,200,000 TRU, with about

20 725,000 TRU representing average prescribed doses.

In one embodiment, an hpHase polypeptide is formulated in a 0.15 M saline solution containing hpHase at a concentration of about 150,000 TRU/cc. The formulation is then injected intravenously at 15,000 TRU/kg body weight of the patient. Alternatively, the enzyme formulation may also be injected

25 subcutaneously to allow the hyaluronidase to perfuse around the tumor site. In a preferred embodiment, hpHase is injected peritumorally or into the tumor mass. In another preferred embodiment, hpHase is formulated as a liposome and is delivered by injection either intravenously or at or near the site of cancerous cells associated with a defect in the LuCa-1 (hpHase) gene. Injection of hpHase

30 intravenously results in hpHase in the tumor site.

Anti-cancer therapy by expression of hpHase in cancerous or pre-cancerous cells

The association of cancer with defects in the human chromosome at 3p21.3 has been described in several tissues (see, e.g.,). The LuCa-1 gene (SEQ ID NO:4) is one of several candidate tumor suppressor genes at 3p21.3. The identity of LuCa-1 and hpHase, as well as the observations that hyaluronidase expression is associated with inhibition of tumor progression and tumor formation in mice (De Maeyer et al. *supra*; Pawlowski et al. *supra*), indicate that the LuCa-1 gene, which encodes hpHase, has activity as a tumor suppressor gene. Therefore, introduction of a nucleotide sequence(s) encoding LuCa-1/hpHase into the genome of cells having a LuCa-1/hpHase defect can repair the defect and thus prevent or inhibit tumor development and/or progression (e.g., replacement gene therapy). The introduced hpHase-encoding sequence can be either constitutively or inducibly expressed upon transformation of the target cell. Alternatively, the LuCa-1/hpHase defect can be repaired by introduction of genetic elements that may enhance expression of hpHase (e.g., introduce an inhibitor of a factor that is responsible for inhibition of hpHase transcription or translation). This latter approach may be useful where the hpHase coding sequence itself is not defective, but rather the defect results from decreased expression of hpHase. For example, transformation of tumor cells or cells in the vicinity of tumor cells (e.g., which can express hpHase which in turn is exposed to the tumor cells) can be used in the treatment of small lung cell carcinoma, squamous lung cell carcinoma, as well as cancers of the breast, ovaries, head and neck, or any other cancer associated with depressed levels of hpHase or with a defective LuCa-1 (hpHase) gene (e.g., a LuCa-1 gene that does not provide for expression of adequate hpHase levels or encodes a defective hpHase that does not provide for an adequate level of hyaluronidase activity) or other defect associated with decreased hpHase activity.

Methods for introduction of a sequence of interest into a host cell to accomplish gene therapy are known in the art. In general, such gene therapy methods include *ex vivo* and *in vivo* methods. *Ex vivo* gene therapy according to the present invention involves, for example, transformation of cells isolated from the patient with an hpHase polypeptide encoding sequence, and implantation of the

transformed cells in the patient to provide a reservoir of hpHase production, preferably at a site within or near the tumor. The implanted cells thus provide secreted hpHase to combat the progression of tumor(s) in the patient. Methods for accomplishing *ex vivo* gene therapy are well known in the art (see, e.g.,

5 Morgan et al. 1987 *Science* 237:1476; Gerrard et al. 1993 *Nat. Genet.* 3:180). *In vivo* gene therapy according to the present invention is used accomplished repair of the hpHase defect within the patient by delivery of hpHase polypeptide-encoding nucleic acid, introduction of the nucleic acid into the cells (e.g., into cancerous cells, pre-cancerous cells, peritumoral cells, or cells capable of

10 expressing and secreting hpHase, preferably into cancerous or pre-cancerous cells), and expression of hpHase by the transformed cells in the patient, thereby repairing the hpHase defect.

Several different methods for transforming cells can be used in accordance with either the *ex vivo* or *in vivo* transfection procedures. For example, various

15 mechanical methods can be used to deliver the genetic material, including the use of fusogenic lipid vesicles (liposomes incorporating cationic lipids such as lipofection; see Felgner et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:7413-7417 (1987)); direct injection of DNA (Wolff, et al., *Science* (1990) 247:1465-1468); pneumatic delivery of DNA-coated gold particles with a device referred to as the

20 gene gun (Yang et al., *Proc. Natl. Acad. Sci. U.S.A.* 1990; 87:1568-9572); and various viral vectors (e.g., non-replicative mutants/variants of adenovirus, retrovirus, adeno-associated virus, herpes simplex virus (HSV), cytomegalovirus (CMV), vaccinia virus, and poliovirus). A review of the different techniques along with a citation of numerous publications in each area is contained within a recent

25 article on human gene therapy (see Morsy et al. 1993 *JAMA* 270:2338-2345).

The formulation to accomplish gene therapy will vary with the gene therapy method used, the route of administration (e.g., administration for systemic gene therapy or administration for transformation of specifically targeted cells), and the site targeted for transformation (e.g., lung, breast, or ovaries). The hpHase

30 polypeptide-encoding nucleic acid sequence may be naked (i.e., not encapsulated), provided as a formulation of DNA and cationic compounds (e.g., dextran sulfate),

or may be contained within liposomes. Alternatively, the DNA of interest can be pneumatically delivered using a "gene gun" and associated techniques which are well known in the art (Fynan et al. 1993 *Proc. Natl. Acad. Sci. USA* 90:11478-11482). The various elements of the construct for hpHase expression in the target
5 cell (e.g., the promoter selected, the presence of elements to enhance expression) will also vary according to the cell type and level of hpHase expression desired.

The amount of DNA administered will vary greatly according to a number of factors including the susceptibility of the target cells to transformation, the size
10 and weight of the subject, the levels of hpHase expression desired, and the condition to be treated. For example, the amount of DNA injected into a human breast tumor is generally from about 1 μ g to 200 mg, preferably from about 100 μ g to 100 mg, more preferably from about 500 μ g to 50 mg, most preferably about 10 mg. Generally, the amounts of DNA for human gene therapy can be
15 extrapolated from the amounts of DNA effective for gene therapy in an animal model. For example, the amount of DNA for gene therapy in a human is roughly 100 times the amount of DNA effective in gene therapy in a rat. The amount of DNA necessary to accomplish cell transformation will decrease with an increase in the efficiency of the transformation method used.

20

Numerous other uses for phase, anti-phase antibodies, and hpHase polypeptide-encoding nucleotide sequences are readily apparent to one of ordinary skill in the art. Nucleotide sequences encoding hpHase polypeptides can be used in hybridization screening methods to detect other hyaluronidases having homology
25 to hpHase.

DEPOSITS

The hybridoma cell lines 17E9 and 4D5, which produces an anti-hpHase antibody that binds native hpHase, has been deposited on behalf of The Regents
30 of the University of California, 300 Lakeside Drive, 22nd Floor, Oakland, California 94612 with the American Type Culture Collection (ATCC), Rockville,

Maryland, U.S.A. for patent purposes. The deposit of the hybridoma cell line 17E9 was received by the ATCC on October 17, 1996, ATCC Designation ATCC HB-12213. The deposit of the hybridoma cell line 4D5 was received by the ATCC on October 17, 1996, ATCC Designation ATCC HB-12214. The hybridoma cells
5 were deposited under the conditions specified by the Budapest Treaty on the international recognition of the deposit of microorganisms (Budapest Treaty).

Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

10

EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to carry out the invention and is not intended to limit the scope of what the inventors regard as their
15 invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

20

Example 1: Hyaluronidase (HAse) Activity Assay Using Covalently-Bound Biotinylated Hyaluronic Acid

Human umbilical cord hyaluronic acid (HA; ICN) was biotinylated through free carboxyl groups with biotin hydrazide (Pierce), coupled using
25 1-ethyl-dimethylaminopropyl carbodiimide (EDC, Sigma). 100 mg of hyaluronic acid (HA) was dissolved in 0.1 M Mes pH 5.0, and biotin hydrazide (Pierce), dissolved in DMSO, was added to a final concentration of 1 mM. EDC was dissolved into the HA/biotin solution to a final concentration of 0.03 mM and left stirring overnight at 4°C. Uncoupled biotin was then removed through exhaustive
30 dialysis against distilled water (dH2O) and was stored at -20°C at a final concentration of 1 mg/ml.

Up to about 10 days before performing the hyaluronidase assay, the biotinylated hyaluronic acid (bHA) was then covalently coupled to Covalink-NH microtiter plates (NUNC, Placerville, NJ) at a final concentration of 5 μg of bHA/well with 9.2 μg /well Sulfo-NHS (Pierce) and 6 μg /well fresh EDC overnight at 4°C. Unbound HA was removed through washing plates with wash buffer (PBS with 2 M NaCl, 50 mM MgSO_4 , 0.05% Tween-20). Incubations for plasma hyaluronidase activity were performed by diluting human plasma 1:400 in assay buffer (0.1 M Formate pH 3.7, 0.1 M NaCl, 1% Triton X-100, 0.02% Azide, 5 mM CaCl_2 and 5 mM sacharrolactone (to inhibit exoglycosidase activity)) and incubating 100 μl samples in the bHA plates for 30 min at 37°C. The reaction was terminated through the addition of 6 M Guanidine HCl and washed three times with PBS wash buffer.

Residual undigested hyaluronic acid remaining covalently bound to the plate was reacted with avidin peroxidase(ABC-kit Vectastain) using o-phenylendiamine as a substrate. Fluorescence associated with bHA was detected using a microtiterplate reader at 492 nm. A standard curve was generated through co-incubation of serial dilutions of bovine testicular hyaluronidase (WYDASE™) and activity of unknown samples were interpolated through a four parameter curve fit to yield values given in relative Turbidity Reducing Units (rTRU)/ml related to the standardized commercial preparations of hyaluronidase.

The bHA hyaluronidase activity assay is approximately 1,000 times more sensitive than the conventional colorimetric assay (Afify et al. *supra*) and requires only 60 min to complete. Moreover, the assay does not require the tedious preparation of a biotinylated-HA binding aggrecan peptide from bovine nasal cartilage as used in conventional ELISA-like assays (Stern et al. *supra*; Delpech et al. *supra*). The assay can also be used to detect Hase activity from cell cultures that produce very low levels of activity by using longer incubation periods.

Example 2: Purification of Human Plasma Hyaluronidase Using the Biochemical Purification Method of the Invention

hpHase was purified in a three-step biochemical procedure: 1) temperature-induced detergent hpHase extraction of human plasma; 2) Fast Flow-S cation
5 exchange chromatography; and 3) hydroxyl-apatite resin. Hyaluronidase activity and protein concentration were determined at each stage of purification, and the specific hyaluronidase activity calculated. Hyaluronidase activity was determined using the assay in Example 1. Protein concentration was determined by both
10 absorbance at 280 nm and with the Biorad (Burlingame, CA) protein microassay kit (Tengblad 1979 *Biochim. Biophys. Acta* 578:281-289) in 96 well plates, using crystallized bovine serum albumin as a standard and read at 595 nM.

1) Temperature-induced detergent phase extraction

Two liters of outdated human plasma, obtained from either UCSF or Irwin Memorial blood donor centers, were routinely used for enzyme purifications. Two
15 liters of chilled human plasma were dissolved in a solution 0.02% Sodium Azide, 50 mM NaCl, 5% sucrose and 7.5% Triton X-114 (Boehringer Mannheim) were dissolved at 4°C with stirring for 90 min followed by centrifugation at 10,000 x g for 30 min to remove insoluble material. The plasma was then subjected to temperature-induced phase extraction at 37°C to separate the detergent-rich and
20 detergent-poor phases. The extract was centrifuged at 10,000 x g for 30 min at 37°C to clarify the two phases. The detergent-rich phase was removed and diluted to 2 liters with cold 50 mM Hepes, pH 7.5, 0.15 M NaCl. The solution was then allowed to mix thoroughly on ice followed by repartitioning at 37°C with centrifugation. This was repeated three times in order to increase the specific
25 activity of the hyaluronidase that partitioned into the detergent phase. The final detergent-rich phase was enriched 60-fold for hpHase specific activity relative to the human plasma starting material as determined by the Hase activity assay of Example 1.

2) Fast Flow-S cation exchange chromatography

The final detergent-rich phase from step 1) was diluted six-fold with 20 ml of equilibrated SP-Sepharose cation exchange resin (Pharmacia) in 25 mM Mes, pH 6.0 and stirred overnight at 4°C. The beads were collected through
5 centrifugation and washed extensively with 25 mM Mes, pH 6.0, containing 46 mM octylglucoside (Boehringer Mannheim). Hyaluronidase was eluted from the beads through the addition of 0.3 M NaCl in Mes pH 6.0 buffer with several washes. The SP-Sepharose eluant was concentrated on a YM3 (Amicon) membrane and desalted into 10 mM PO₄ pH 7.4 with 25 mM NaCl, 46 mM
10 octylglucoside on a f.p.l.c. Fast-Desalting column.

3) Hydroxyl-apatite resin

The hyaluronidase preparation from step 2) was combined with 10 ml of equilibrated hydroxylapatite resin (Biorad) and left on a rocker overnight at 4°C. hpHase did not adsorb to the resin under these conditions and was recovered in
15 the supernatant. hpHase recovered in the supernatant was purified to homogeneity as determined by electrophoretic analysis on silver-stained 12.5% polyacrylamide gels on a Pharmacia Phast Gel System.

Table 1 is a summary of the hyaluronidase activity, protein concentration,
20 and specific activities of the hpHase-containing fractions during each of the purification steps described above.

Table 1. Biochemical Purification of hpHase

Purification Step	Volume (ml)	Activity (rTRU/ml)	Protein (mg/ml)	Specific Activity (rTRU/mg)	X-Fold Purification
Starting Material (human plasma)	2,100	35	86	0.406	1.0
1) Final Detergent Phase	650	33	1.3	25.4	63
2) Fast Flow-S Cation Exchange	60	298	0.85	350.6	875
3) Hydroxyl-Apatite (unbound)	15	907	0.0015	6.0×10^5	1.5×10^6

Summary of results of biochemical hpHase purification

Purification of hpHase using the biochemical method described above revealed that hpHase has the unique property of partitioning into the temperature-induced detergent phase. hpHase partitioning into the detergent-rich phase is usually observed with certain integral membrane or lipid-anchored proteins (Bordier 1981 *J. Biol. chem.* 256:1604-7). Treatment of hpHase with phosphatidylinositol specific phospholipase-C (which cleaves glycosyl-phosphatidyl inositol (GPI) anchors), phospholipase-D (which cleaves GPI anchors), or N-glycosidase-F (which cleaves N-linked glycosylated moieties) did not alter the hpHase partitioning properties of hpHase (although N-glycosidase-F treatment combined with gel zymography did show that hpHase has at least two N-linked glycosylation sites).

These experiments suggested that hpHase is an integral membrane protein or has a phospholipase-resistant anchor, similar to the GPI anchor of human erythrocyte acetylcholinesterase (Roberts et al. 1988 *J. Biol. Chem.* 263:18766-75). Although GPI anchors are normally associated with proteins that reside on the extracellular domain of the plasma membrane, there are examples of GPI anchored

proteins in serum (e.g., CD59, Vakeva et al. 1994 *Immunology* 82:28-33). Furthermore, several of the neutral PH20 sperm hyaluronidases have GPI-specific phospholipase-C-susceptible anchors (Gmachl et al. 1993 *FEBS Lett.* 336:545-8; Thaler et al. 1995 *Biochemistry* 34:7788-95), but no GPI-anchor-like post-translation modifications have been described in an acid active hyaluronidases. Moreover, such post-translational modification would not be obvious from the amino acid or nucleotide sequence encoding such acid active hyaluronidases (i.e., no specific sites for post-translational modification by addition of a fatty acid or GPI anchor have been described). For example, there is no presently known consensus sequence for GPI-anchored proteins. The consensus sequence for isoprenylated proteins (CAAX; SEQ ID NO:10) is known; however, hpHase does not contain this sequence.

hpHase displayed other strongly amphiphilic characteristics. For example, hpHase precipitated from solution when the ionic strength of the solution in which hpHase was dissolved was lowered through dialysis. hpHase was enriched within the 0-30% ammonium sulfate fraction. The concavalin-A binding properties of hpHase revealed that hpHase is a mannose-containing glycoprotein.

Gel filtration chromatography on an S200-f.p.l.c. column (Pharmacia) at neutral pH resulted in elution of the Hase activity peak within the void volume at approximately 120 kDa. When gel filtration chromatography was performed in the absence of non-ionic detergents, the Hase activity peak eluted at approximately 60 kDa. This difference in the molecular weight associated with the hpHase activity is likely due to hpHase aggregation or oligimerzation (e.g., through intramolecular bonds) in the presence of detergents. hpHase is very stable at 37°C in the presence of non-ionic detergents; thus, Triton X-114 phase extractions are ideal as an initial fractionation step, particularly since few plasma proteins partition into the detergent-rich phase.

During cation exchange chromatography to remove detergent and further purify hpHase, it was found that the addition of non-ionic detergents and maintenance of at least 50 mM NaCl was essential to prevent precipitation of the enzyme. Batch adsorption chromatography preserved activity in comparison to

column chromatography. Interestingly, hpHase activity was not retained when concentrating hpHase-containing preparations on 10 kDa cut off membrane in the presence of 60 mM octylglucoside; however, hpHase activity was retained by using a 3 kDa cutoff membrane. These results suggest that the Stokes radius for
5 hpHase deviates substantially from a spherical model. These data show that hpHase does not have a spherical shape in the presence of octyl glucoside detergent as assumed for most globular proteins, but rather likely has more rod-like shape. hpHase in one dimension appears to have a molecular weight of 6-7kDa protein rather than a 57kDa protein. The protein can thus in one dimension pass
10 through barriers reserved to small proteins and peptides, which would explain the finding of the plasma enzyme in human urine (see *infra*, usually the kidney excludes proteins <50-60kDa). Thus, patients may excrete a significant amount of hpHase into their urine if hpHase is not incorporated into liposome or liposome-like structures.

15 The post SP-Sepharose preparation of hpHase was purified to homogeneity through adsorption of contaminating proteins with hydroxyl apatite resin, resulting in an overall purification of 1.5 million fold. SDS-PAGE electrophoresis and silver staining of an hpHase sample from this final purification step revealed a single band, thus indicating that hpHase had been purified to electrophoretic
20 homogeneity. The specific activity of the purified enzyme (600,000 rTRU/mg as tested using the assay described in Example 1) was roughly 6-fold that of the reported values for the sperm hyaluronidase, PH20. The purified enzyme migrated on SDS electrophoresis gels with a relative molecular mass of 57 kDa, though levels of nonionic detergents in the final preparation made molecular mass analysis
25 variable.

Example 3: Screening Assay for Anti-Native Acid Active Hase (aaHase) Antibodies

Biotinylated HA (bHA) was prepared as described in Example 1.
30 Approximately 5 μ g of bHA/well and 1.25 μ g/well of goat anti-mouse IgG (Jackson Immunolabs) were covalently coupled to Covalink-NH microtiter plates

(NUNC, Placerville, NJ) with 9.2 μ g/well Sulfo-NHS (Pierce) and 6 μ g/well fresh EDC overnight at 4°C. Unbound HA and goat antibody were removed through washing plates with wash buffer (PBS with 2M NaCl, 50mM MgSO₄, 0.05% Tween-20).

- 5 Anti-native acid active Hase (anti-native aaHase) antibodies were screened by incubating candidate antibodies with a sample containing an acid active Hase in a neutral pH buffer composed of 1% Triton X-100 in phosphate-buffered saline (PBS). If the sample was a diluted sample (e.g., a diluted plasma sample), the neutral pH buffer additionally contained 5 mg/ml bovine serum albumin (BSA).
- 10 The aaHase-containing sample was incubated with the candidate antibodies to allow for formation of native aaHase-antibody complexes. The sample was then placed into bHA/anti-mouse antibody wells and incubated to allow for formation of anti-mouse antibody/anti-native aaHase/aaHase antibodies in the neutral pH buffer. A sample containing antibodies that do not bind aaHase can be used as a
- 15 negative control.

- Detection of antibodies bound to native aaHase was accomplished by first washing away unbound material with neutral pH buffer, then replacing the neutral pH buffer with the acidic assay buffer used in the Hase activity assay described in Example 1 (0.1 M Formate pH 3.7, 0.1 M NaCl, 1% Triton X-100, 0.02%
- 20 Azide, 5 mM CaCl₂ and 5mM sacharrolactone). The shift in the pH from neutral to acidic allows for the Hase activity of any bound aaHase to degrade the covalently bound bHA. The reaction was terminated by addition of 6 M guanidine HCL and washed three times with PBS wash buffer (PBS, 2 M NaCl, 50 mM MgSO₄, 0.05% Tween-20). Degradation of bHA was detected by reaction of the
- 25 wells with avidin peroxidase using o-phenyldiamine as a substrate and reading the plates at 492 nm as described in Example 1. Anti-native aaHase antibodies were identified by degradation of bHA. The assay takes advantage of the aaHase characteristic that aaHases have no affinity for the HA substrate above pH 4.5 (as determined through HA-Sepharose affinity chromatography with hpHase, data not
- 30 shown).

Example 4: Generation and Identification of Anti-hpHase Monoclonal Antibodies

hpHase isolated from the post-hydroxyl apatite step in the purification method of Example 2 was used to immunize five 6 week old female Balb-c mice according to methods well known in the art (see, e.g., Harlow and Lane, *supra*).
5 Briefly, the hpHase was combined with Freund's complete adjuvant and injected intraperitoneally into the mice. At 21 day intervals, the animals were boosted with protein plus Freund's incomplete adjuvant. The fourth and final boost contained hpHase in Freund's incomplete adjuvant was injected intravenously. Serum samples were obtained, the mice sacrificed, and spleen cells from two of the five
10 isolated for preparation of hybridoma cell lines according to methods well known in the art (see, e.g., Harlow and Lane, *supra*; Schrier et al. *supra*). Fusion of the balb/c cells was performed with sp2/0 myelomas.

Antibody-secreting hybridomas were screened using the anti-native aaHase antibody assay of the invention as described Example 3. Briefly, hybridoma
15 supernatants from 20 fusion plates were incubated with diluted human plasma for 60 min at 37°C followed by incubation in the bHA/anti-mouse-IgG plates for 60 min at 37°C. The plates were washed five times with PBS containing 1% Triton X-100 and 10 mg/ml BSA to remove non-immunoprecipitated hyaluronidase. The acidic formate assay buffer (pH 3.7) was then added to the
20 wells and incubated at 37°C for 60 min. After stopping the reaction with 6 M guanidine, undegraded bHA remaining in the wells was detected by reaction of the wells with avidin peroxidase using o-phenyldiamine as a substrate and reading the plates at 492 nm as described in Example 3.

Eight clones were identified from the original 20 hybridoma fusion plates.
25 Two clones, 17E9 (producing IgG_{2a} class antibody; kappa chain) and 4D5 (producing IgG₁ class antibody; kappa chain), were used to generate ascites. Immunoglobulin from ascites of single-cell cloned hybridoma lines were generated in Balb/c mice and purified through protein-A affinity chromatography. Neither the 17E9 or 4D5 antibodies blocked hpHase activity. The 17E9 antibody is
30 preferred for use in immunoaffinity purification and immunoprecipitation; the 4D5 antibody is preferred for use in immunohistochemistry.

Example 5: Immunoprecipitation of hpHase using the 17E9 Monoclonal Antibody

hpHase was immunoprecipitated from human plasma using either purified IgG2a from the 17E9 anti-native hpHase clone or a control IgG2a non-specific antibody. Human plasma was diluted in R.I.P.A. buffer (1% NP40, 1% Deoxycholate, 1% Triton X-100, 5 mM EDTA in PBS). Protein-A Sepharose conjugated to serial dilutions of either control IgG2a antibody or 17E9 antibody was added to the sample. Unbound material was separated from the beads by centrifugation and the Hase activity remaining in the supernatant detected using the assay described in Example 1. As shown in Fig. 4, immunoprecipitation with the 17E9 antibody resulted in removal of essentially all of the detectable acid active hyaluronidase activity. Further, the 17E9 antibody did not bind bovine testicular hyaluronidase (PH20), suggesting that 17E9 does not bind a carbohydrate moiety or other moiety that may be shared by PH20 and hpHase.

Example 6: Immunoaffinity Purification of hpHase

hpHase was purified to homogeneity from raw human plasma in a single step using the 17E9 antibody bound to an immunoaffinity chromatography column. Approximately 3 mg of purified IgG from the 17E9 hybridoma clone ascites was coupled to a 1 ml High Trap-NHS activated column (Pharmacia) as per the manufacturer's instructions. Human plasma was diluted 1:2 with R.I.P.A. buffer, and passed over the 17E9 IgG antibody column. The column was then washed with PBS containing 2M NaCl, 100 mM octylglucoside followed by washing with 100 mM citrate pH 4.0. hpHase was eluted with 100mM citrate, pH 3.0, with 150 mM NaCl. hpHase eluted as a sharp, homogenous peak at pH 3.0. The specific activity of immunoaffinity purified hpHase ranged from about 4×10^5 rTRU/mg. Thus very little, if any, activity is lost in the immunoaffinity purification process relative to the biochemical purification process. Moreover, immunoaffinity purification resulted in a yield of hpHase approximately 100% of the hpHase in the sample; Hase assay detected no Hase activity remaining in the unbound fraction.

Example 7: Amino Acid Sequencing of hpHase

Approximately 50 μ g of immunoaffinity purified hpHase from Example 6 was digested with Lys-C (to generate N-terminal fragments) or trypsin (to generate internal fragments) and chromatographed on a Vydac C-18 reverse phase column to separate fragments. Purified peptides were sequenced on a gas phase Edman sequencer. For N-terminal amino acid sequencing, the purified protein was immobilized on a Prosorb membrane (ABI). N-terminal amino acid sequencing was accomplished by automated protein sequence analysis using gas phase Edman degradation. Fasta queries (Pearson et al. 1988 *Proc. Natl. Acad. Sci. USA* 85:2444-8) of N-terminal and 5 separate internal amino acid sequence derived from the Lys-C digests of purified hpHase (SEQ ID NO:1) revealed identity with the predicted amino acid sequence of the LuCa-1 gene (SEQ ID NO:3; GenBank accession no. u03056).

Example 8: Characterization of LuCa-1/hpHase

As described in the Example 7 above, hpHase is identical to LuCa-1. The LuCa-1 gene is one of a group of 17 sequences located on chromosome 3p.21.3 that display a 100% loss of heterozygosity in small cell lung carcinoma lines Dietrich et al. 1966 *Clin. Chim. Acta* 13:746-52). 3p.21 deletions spanning the region containing LuCa-1 have also been described in non-small cell lung carcinomas, mammary carcinoma, carcinoma of the prostate and head and neck carcinoma. Chromatofocusing of hpHase on a MONO-P f.p.l.c. system provided additional data regarding the identity of hpHase (SEQ ID NO:1) and LuCa-1 (SEQ ID NO:3). hpHase eluted at pH 6.5, very close to the LuCa-1's calculated theoretical isoelectric point of 6.58.

Pearson Lipman alignment (Pearson et al. *supra*) of LuCa-1 (SEQ ID NO:3) and human testicular hyaluronidase PH20 (SEQ ID NO:11) amino acid sequences revealed that LuCa-1 and PH20 share over 40% sequence identity and 60% percent homology (Fig. 5). PH20 is a predominantly sperm-specific neutral hyaluronidase and shares considerable homology with the venom hyaluronidases found in bee and yellow jacket vespids venoms (Gmachl et al. 1993 *Proc. Natl. Acad. Sci. USA*

90:3569-73). The striking homology between a strictly acid active plasma hyaluronidase (LuCa-1/hpHase) and PH20 is quite surprising given their very different pH optima, suggesting that all mammalian β ,1-4 hyaluronidases, both neutral and acidic, are in members of a highly conserved family of enzymes.

5 Hydropathy plots of LuCa-1/hpHase (Fig. 6) did not reveal any rich hydrophobic domains (e.g., such as those associated with integral membrane proteins) that would explain the phase partitioning properties of hpHase described in Example 2. These data thus suggest that partitioning of LuCa-1/hpHase into the detergent-rich phase is due to a post-translational modification (e.g., fatty acid
10 chain modification or phospholipase C/phospholipase D/N-glycosidase F-resistant GPI anchor.

Example 9: Isolation of LuCa-1/hpHase-encoding DNA

cDNA encoding the LuCa-1 coding sequence was isolated using two rounds
15 of nested PCR reactions with a λ gt10 5'-STRETCH PLUS Human Liver cDNA library (Clontech Laboratories, Inc., Palo Alto, California, USA). In the first round of PCR, which amplified nucleotides 590 to 1948 of the LuCa-1 cDNA, the following primers were used: LuCaF1 (5'- CAGGTTGTCCTGCACCAGTC
-3')(SEQ ID NO:5), and LuCaR1 (5'- ATGTGCAACTCAGTGTGTGGC-3')(SEQ
20 ID NO:6). The PCR reaction was performed in a total reaction volume of 50 μ l containing 1 μ l of liver cDNA library, 1 μ l each of 25 mM primers, 1 μ l of 10 mM dNTPs (Gibco BRL, Grand Island, New York, USA), 2 units of Pfu DNA polymerase (Stratagene Cloning Systems, La Jolla, California, USA), buffered with 10 mM Tris-HCl, 50 mM KCl, and 2 mM $MgCl_2$. The PCR reaction conditions
25 included a 1 min denaturation step at 95°C, 1 min annealing step at 60°C, and a 1 min extension at 74°C for 37 cycles, followed by a final 7 min extension at 74°C.

A second round of PCR amplification was used to amplify nucleotides 612 to 1925 of the Luca-1-encoding cDNA with the nested PCR primers: LuCaF2 (5'-
30 GTGCCATGGCAGGCCACC -3')(SEQ ID NO:7) and LuCaR2 (5'- ATCACCACATGCTCTTCCGC -3')(SEQ ID NO:8). Primer LuCaF2 was

phosphorylated prior to the PCR reaction in order to facilitate subsequent unidirectional cloning of the PCR product. Phosphorylation was carried out by incubating 2 ml of 25 mM LuCaF2 with 10 units of T4 Polynucleotide Kinase, 1 ml of 10 mM ATP buffered in 50 mM Tris-HCl pH 7.5, 10 mM MgCl₂, 5 mM dithiothreitol, and 0.1 mM spermidine in a final volume of 9 ml. The reaction was carried out in a total volume of 50 ml containing 1 ml of the final product from the first PCR reaction as template, 4 ml of phosphorylated LuCaF2, 1 ml of 25 mM LuCaR2, 1 ml 10 mM dNTPs (Gibco BRL), buffered with 10 mM Tris-HCl, 50 mM KCl, and 1.5 mM MgCl₂ and containing 0.5 units of Taq DNA polymerase (Gibco BRL). The second PCR reaction consisted of 7 cycles of a 1 min denaturation at 95 °C, a 1 min annealing at 58°C, and a 1 min extension at 72°C, followed by a final extension at 72°C for 7 min.

The product of the PCR reaction was ligated into 2 µl of the unidirectional TA-expression vector pCR3.1-Uni (Invitrogen, San Diego, CA, USA) with T4 DNA ligase. The ligated vector was used to transform One Shot™ TOP10F' (Invitrogen) competent cells, which were plated out onto LB agar plates containing 50 µg/ml ampicillin and incubated overnight at 37°C. Colonies were tested for the presence of LuCa-1 inserts by growing overnight colonies in 10 ml LB broth with 50 µg/ml ampicillin, and by plasmid purification with the Wizard™ Plus Miniprep DNA Purification System (Promega Corporation, Madison, WI, USA), followed by restriction mapping of the vector with Dra III endonuclease (Boehringer Mannheim, Indianapolis, IN, USA) and automated fluorescent sequencing on an ABI Prism™ DNA sequencer.

Restriction mapping with DraIII and sequencing of the plasmid showed that the cloned sequence was identical to the LuCa-1 sequence available in the GenBank database except for a Val₂₇-Leu substitution found in the N-terminus. Sequencing of the active PCR product from the liver λgt10 cDNA library revealed a G-C discrepancy between the sequence obtained and the published LuCa-1 sequence at the position corresponding to the 27th amino acid residue. Thus, either the published Luca-1 sequence contains an error or the clone sequenced here is an allelic variant.

Example 10: Expression of recombinant LuCa-1 in Cos-7 cells

The purified LuCa-1-encoding cDNA in the pCR3.1-Uni expression plasmid was transfected into 50% confluent cos-7 cells in T75 flasks for five hours using 9 μ g of DNA with 60 μ l of Lipofectamine in 20 ml of DME/F12 50/50 mix with
5 non-essential amino acids (UCSF cells culture facility) containing insulin, transferrin and selenium (Gibco BRL). The transfected cells were then grown for an additional 48 hr in DME/F12 50/50 mix containing 10% fetal bovine serum.

10 Example 11: Purification and Characterization of Recombinant LuCa-1 Expressed in Cos-7 Cells

Recombinant LuCa-1 was further characterized to further verify that LuCa-1 is identical to hpHase by virtue of its ability to bind anti-native hpHase antibodies and exhibit acid active Hase activity. Recombinant LuCa-1 was expressed in cos-
15 7 cells as described above and subjected to the following experiments: 1) Hase activity assays using anti-hpHase antibodies; 2) immunoprecipitation of LuCa-1 using anti-native hpHase antibodies; 3) gel zymography of anti-native hpHase antibody-precipitated LuCa-1; and 4) determination of the pH optima of the acid active Hase activity associated with LuCa-1. Cells transfected with 9 μ g of the
20 pCR3.1-Uni vector containing the chloramphenicol transferase gene served as a negative control (mock-transfected cells).

Hase activity of recombinant LuCa-1 using anti-hpHase antibodies in the enzyme capture assay

Cos-7 cells expressing hpHase and their conditioned media were separately
25 extracted with 2% Triton X-114 followed by temperature-induced detergent phase separation as described in Example 2. The detergent-rich phase extracts were analyzed for Hase activity using the enzyme immuno-capture assay of Example 1. Triton X-114 detergent phase extracts of both the cell layer and conditioned media contained an acid active Hase activity as detected in the assay of the invention
30 (Fig. 7A). Mock-transfected control cells secreted a detectable, low-level, acid active Hase activity (Fig. 7A).

To further assess the Hase activity of recombinant hpHase, HEK 293 cells were stably transfected with hpHase-encoding DNA in a manner similar to that described above for transformation of Cos-7 cells. Briefly, the hpHase-encoding construct was transfected into HEK 293 cells using 9 μ g of DNA with 60 μ l of Lipofectin. After 48 hrs, the cells were plated out using the limiting dilution method into 24 well plates with 500 μ g/ml G418. After 14 days, the conditions media of resistant colonies was assayed for hyaluronidase activity as described above. Colonies with high level expression were expanded for further characterization. Analysis of the recombinant hpHase was carried out by growing an HEK-293 cell line overexpressing the hpHase-encoding construct for 48 hrs in serum free media. The condition media was passed through a 17E9 anti-hpHase immunoaffintiy column. Recombinant hpHase was eluted using the protocol above. As shown in Fig. 7B, both the cell layer and conditioned media contained an acid active hpHase activity; mock-transfected cells secreted no detectable hyaluronidase activity.

These data show that LuCa-1 exhibits an acid active Hase activity, thus lending further support to the observation that hpHase and LuCa-1 are identical.

Immunoprecipitation of recombinant LuCa-1 with anti-hpHase antibodies

Binding of anti-native hpHase antibodies to LuCa-1 was tested in the immunoprecipitation method described in Example 5 using the 17E9 anti-hpHase monoclonal antibody described in Example 4 bound to protein A Sepharose. The 17E9 antibodies immunoprecipitated LuCa-1 from both the LuCa-1 expressing-cells and from the conditioned media. However, the acid active Hase activity of mock transfected cells was not immunoprecipitated with the 17E9 antibody-protein A Sepharose beads. These experiments show that hpHase and LuCa-1 share the antigenic epitope bound by the 17E9 anti-native hpHase antibody. Moreover, this epitope is unique to hpHase and LuCa-1, since the 17E9 did not bind an acid active Hase expressed by mock transfected cos-7 cells.

Gel zymography of LuCa-1 immunoprecipitated with anti-hpHase antibodies

Recombinant LuCa-1 was immunoprecipitated with the 17E9 anti-native hpHase antibody was also tested for acid active Hase activity in substrate gel zymography experiments according to methods known in the art (Afify et al. *supra*). Briefly, cell lysates and conditioned media from LuCa-1-transfected cells (test samples) and mock transfected cells (chloramphenicol-transferase encoding plasmid; negative control) were immunoprecipitated with anti-native hpHase bound to protein A Sepharose beads. The samples were suspended in SDS sample buffer and electrophoresed in 10% polyacrylamide gels containing 40 μ g/ml HA. Samples containing immunoprecipitated hpHase (Example 5) and bovine testicular hyaluronidase (WYDASE™ 150 rTRU/ml) served as positive controls. Incubations were performed as described by Afify et al. (*supra*) and digested HA detected by staining with Alcian blue/Acetic acid followed by Commassie staining to enhance the Alcian/carbohydrate stain and visualize protein.

The region of clearing the SDS-HA gel corresponded with the same relative molecular mass as the immunoaffinity purified hpHase preparation. No zone of clearing, and thus no immunoprecipitated acid active hyaluronidase activity was observed in the mock-transfected cell samples.

pH optima of Hase activity of recombinant LuCa-1

The Hase activity assay described in Example 1 was used to determine the pH optimum of the recombinant LuCa-1 Hase activity. The Hase activity assay of Example 1 was performed as described, except that the assay buffer was composed of 0.1 M Formate pH 3-4.5, Acetate pH 5.0, Mes pH 6.0, Hepes pH 7.0-8.0, 0.1 M NaCl, 1% Triton X-100, 0.02% Azide, 5 mM CaCl₂ and 5 mM sacharrolactone. Sigma Type VI-S testicular hyaluronidase (3,000 TRU/mg solid), which contains the neutral Hase PH20 (maximal Hase activity at about pH 7.5), was used as a comparison. The Type VI-S sample was treated identically, except that the assay buffer for this sample was prepared without NaCl (NaCl inhibits neutral Hase activity). As shown in Fig. 8, the pH optima curve of recombinant LuCa-1 displays the same strictly acid active profile as immunoaffinity purified

hpHase. Neither LuCa-1 nor hpHase display Hase activity above pH 4.5; in contrast, Type VI-S testicular hyaluronidase exhibited Hase activity only above pH 7.5.

In summary, biochemical, molecular, and immunological criterion strongly
5 indicate that LuCa-1 and hpHase are identical.

Example 12: Organ Survey of LuCa-1/hpHase and Transient Expression of LuCa-1/hpHase

The primers described in Example 11 that were used to amplify the 1.3kb
10 coding region of the LuCa-1 cDNA were used to analyze the tissue distribution of LuCa-1/hpHase transcripts according to methods well known in the art. Amplified PCR products from λ gt10 cDNA libraries of various tissues were detected in heart, kidney, liver, lung, placenta, and skeletal muscle, but were not detected in brain. Heart tissue exhibited one of the highest levels of LuCa-1/hpHase transcript
15 production.

Following procedures similar to those described above, other hyaluronidases having substantially the same sequence as hpHase of the invention can be purified, cloned, and expressed.

20 Example 13: Biochemical Purification of a Form of hpHase from Urine

The urine form of hpHase was purified from human urine according to the biochemical purification method described in Example 2, except that concentrated urine was used as the sample in lieu of outdated human plasma. Urine hpHase partitioned into the detergent-rich Triton X-114 phase in a manner similar to that
25 of hpHase, suggesting the urine hpHase contains a lipid modification similar to that of hpHase. The isoelectric point of urine hpHase is 6.5 as determined by elution in chromatofocusing on Mono-P f.p.l.c. Human urine hpHase immunoprecipitated with the anti-native hpHase antibody 17E9.

Gel zymography revealed two bands of Hase activity in crude plasma and
30 urine samples. In plasma, Hase activity was detected in two bands corresponding to 57 kDa and 46-47 kDa; the 57 kDa band was the predominant species. In urine,

Hase activity was also detected in two bands corresponding 57 kDa and 46-47 kDa; however, neither species was predominant (i.e., the bands appeared in the urine sample with equal intensity). These data suggest that hpHase and urine hpHase are present in plasma and urine, respectively, in two distinct modified forms (e.g., as a holoprotein and a proteolytic or otherwise modified fragment), or that there are two distinct acid active Hase proteins present in urine and in plasma.

Example 14: Expression of hpHase in Metastasis-Derived Carcinomas

Several carcinoma lines (SCC 10A, SCC 10B, HSC-3, NIC H740, and DMS 153) were examined for the production of human plasma hyaluronidase by characterizing the levels of 17E9 immunoreactive hpHase activity using the immunoprecipitation assay described in Example 5 above. hpHase levels were examined in both the conditioned media and in the cell layer itself. Normal foreskin human keratinocytes served as a positive control. The SCC 10 A cell line is derived from a primary tumor of a laryngeal carcinoma; the SCC 10B cell line is from a lymph node metastasis from the same patient. The HSC-3 cell line is derived from a lymph node metastasis. The NIC H740 cell line is a small cell lung carcinoma containing a homozygous deletion of the region of chromosome 3p21.3 encoding hpHase. The DMS 153 cell line is a classic small lung cell carcinoma cell line (as opposed to a small lung cell carcinoma variant cell line).

As shown in Fig. 9, the SCC 10A cell lines produces hpHase activity levels comparable to that of normal human keratinocytes, but the levels of activity in the lymph node metastasis from the same patient (SCC 10B) is completely absent in the cell layer or conditioned media. The HSC-3 derived cell line exhibited no detectable hpHase activity (the hpHase activity assay used can accurately detect at least 1/100th of the activity found in normal keratinocytes). Neither of the small cell lung carcinoma lines NIC H740 or DMS 153 have detectable activity. NIC H740 and DMS 153 cells also did not generate any detectable hpHase-encoding transcripts (data not shown). These data suggest that loss of hpHase expression is strongly correlated with tumorigenesis. The present

inventors have identified no metastasis-derived carcinoma that produces a functional plasma hyaluronidase gene product. Some non-metastatic lines derived from primary carcinomas do have functional activity, suggesting that complete loss of enzyme function is associated with tumor metastasis.

5

Example 14: Recombinant hpHase Expression System Having Improved Yield

A recombinant hpHase expression system to provide high yields of hpHase in the culture supernatant was developed. hpHase-encoding DNA was operably inserted into the CMV-promoter driven PCR3.1 Uni vector (Invitrogen). This
10 hpHase-encoding construct was then used to transform HEK cells according to methods well known in the art. The HEK cells were then grown in a T225 cm² flask at 37°C, 5% CO₂ in DME H21 (4.5 g/l glucose), 10% FBS media. Conditioned culture media was collected from the cells and the levels of hpHase assessed.

15 While there are often inherent difficulties generating recombinant plasma products with competitive yields to raw plasma, the recombinant hpHase expression system described herein produces over 30-fold the amount of hyaluronidase present in raw human plasma Cohn fraction-I paste (prepared by cold alcohol precipitation according to methods well known in the art) (Table 1). In
20 comparison, transient expression of hpHase in Cos-7 cells expressed about 10% of the levels achieved with HEK cells. Without being held to theory, mammalian cell lines that are adenovirus transformed

Table 1 Levels of hpHase in various preparations

Enzyme Source	Hyaluronidase Activity
Human Plasma	3-6 rTRU/ml
Recombinant Plasma HA'se Conditioned Media	150.0 rTRU/ml

The recombinant hpHase from the mammalian expression system is indistinguishable from the biochemically purified enzyme as determined by using several criteria. Immunoaffinity purified recombinant hpHase migrates with the same molecular mass as biochemically purified hpHase. The amino acid sequence of immunoaffinity purified recombinant hpHase has the same processed N-terminus as the native plasma enzyme (FRGPLLV)(SEQ ID NO:9). Moreover, N terminal sequencing indicates that recombinant hpHase is processed properly in our expression system. Recombinant hpHase has a specific activity equivalent to biochemically purified hpHase as determined by gel zymography. 10U samples of recombinant hpHase, biochemically purified hpHase, and the "purest" commercial preparation of testicular hyaluronidase can depolymerize 100 μ g of high molecular weight HA in 10 min. In addition, recombinant hpHase displays the same detergent-phase partitioning properties as the biochemically purified hpHase, suggesting that the expression system facilitates the correct post-translation processing of recombinant hpHase.

Example 15: Activity of hpHase in Inhibition of Tumor Growth in an Animal Model

The ability of recombinant hpHase to inhibit tumor growth was examined in the HSC-3 head and neck squamous cell carcinoma model, an orthotopic tumor xenografts (transplanted tumors into organ/tissue of origin). Briefly, Nu/Nu mice were orthotopically implanted with 5×10^6 HSC-3 human squamous cell carcinoma cells in Matrigel into the floor of the mouth. The cells were allowed to grow for 7 days. Animals were then segregated randomly into two groups.

Recombinant hpHase was prepared from HEK-conditioned media overexpressing hyal-1 under a CMV promoter using an immunoaffinity column having the 17E9 IgG2a antibody coupled to NHS sepharose. Eluted hpHase was exchanged into bovine HDL complex (Sigma) and dialyzed against Biobeads. HDL complex carrier without hpHase served as a negative control. The material was injected into the mice peritumorally (approximately 10 mm from the primary tumor site) in a 100 μ l volume. Approximately 400 rTRU hpHase was administered per

injection. Injections were repeated every 48 hours. Tumor volume was measured in 3 dimensions using a spherical model. As shown in Fig. 10, administration of hpHase significantly reduced tumor volume.

5 Example 16: Inhibition of Tumor Growth by Expression of hpHase-Encoding DNA in an Animal Model

 The effect of expression of hpHase in tumor cells upon tumor growth was examined by introducing a functional hpHase gene into a carcinoma line, where the hpHase-encoding sequence is under control of an inducible promoter. The
10 steroid-responsive expression plasmid ecdysone was chosen for this study since only those mammalian cells that contain the appropriate steroid-responsive plasmid responded to injection of the stimulating hormone (muristerone). This system thus permits examination of the effects of specific reactivation of the hpHase-encoding gene within the tumor cells and assessment of *in vivo* effects.

15 HSC-3 oral squamous cell carcinoma cells were transfected with Lipofectin (Gibco) at 60 μ l lipofectin, 9 μ g. plasmid per T75 flask with a muristerone inducible gene construct (Ecdysone system, Invitrogen) containing the ecdysone receptor plasmid and encoding hpHase (under inducible steroid control), as well as G418S and zeocin resistance. Cells transformed with the muristerone inducible
20 construct without hpHase-encoding DNA served as a control. The transformed cells were injected in the floor of the mouth of nu/nu mice at 5×10^6 cells per injection as described in Example 15. After 7 days of tumor growth, the mice received an intraperitoneal injection of muristerone (5 mg) every 72 hours to induce expression of the constructs. Tumor volumes were measured as described
25 above in Example 15. In addition, the time to cachexia (defined by a drop in animal mass 15% below starting weight). Survival curves were generated using a Kaplan Meyer analysis using percent cachexic rather than percent surviving.

 As shown in Fig. 11, mice having HSC-3 cells transformed with the hpHase-encoding construct were significantly reduced in tumor growth relative to
30 control mice. Moreover, significantly fewer mice bearing the hpHase-expressing tumor cells progressed to cachexia over the course of 39 days.

The invention now being fully described, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto without departing from the spirit or scope of the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: The Regents of the University of California
- (ii) TITLE OF INVENTION: Human Plasma Hyaluronidase
- (iii) NUMBER OF SEQUENCES: 10
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Robbins, Berliner & Carson, LLP
 - (B) STREET: 201 N. Figueroa Street, 5th Floor
 - (C) CITY: Los Angeles
 - (D) STATE: CA
 - (E) COUNTRY: USA
 - (F) ZIP: 90012-2628
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (vii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Berliner, Robert
 - (B) REGISTRATION NUMBER: 20,121
 - (C) REFERENCE/DOCKET NUMBER: 5555-458C1-XPC
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (213) 977-1001
 - (B) TELEFAX: (213) 977-1003

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 435 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met	Ala	Gly	His	Leu	Leu	Pro	Ile	Cys	Ala	Leu	Phe	Leu	Thr	Leu	Leu
1				5					10					15	
Asp	Met	Ala	Gln	Gly	Phe	Arg	Gly	Pro	Leu	Leu	Pro	Asn	Arg	Pro	Phe
	20							25					30		
Thr	Thr	Val	Trp	Asn	Ala	Asn	Thr	Gln	Trp	Cys	Leu	Glu	Arg	His	Gly
	35						40					45			
Val	Asp	Val	Asp	Val	Ser	Val	Phe	Asp	Val	Val	Ala	Asn	Pro	Gly	Gln
	50					55					60				
Thr	Phe	Arg	Gly	Pro	Asp	Met	Thr	Ile	Phe	Tyr	Ser	Ser	Gln	Leu	Gly
	65				70				75				80		
Thr	Tyr	Pro	Tyr	Tyr	Thr	Pro	Thr	Gly	Glu	Pro	Val	Phe	Gly	Gly	Leu
				85					90					95	

69

Pro Gln Asn Ala Ser Leu Ile Ala His Leu Ala Arg Thr Phe Gln Asp
 100 105 110
 Ile Leu Ala Ala Ile Pro Ala Pro Asp Phe Ser Gly Leu Ala Val Ile
 115 120 125
 Asp Trp Glu Ala Trp Arg Pro Arg Trp Ala Phe Asn Trp Asp Thr Lys
 130 135 140
 Asp Ile Tyr Arg Gln Arg Ser Arg Ala Leu Val Gln Ala Gln His Pro
 145 150 155 160
 Asp Trp Pro Ala Pro Gln Val Glu Ala Val Ala Gln Asp Gln Phe Gln
 165 170 175
 Gly Ala Ala Arg Ala Trp Met Ala Gly Thr Leu Gln Leu Gly Gly Ala
 180 185 190
 Leu Arg Pro Arg Gly Leu Trp Gly Phe Tyr Gly Phe Pro Asp Cys Tyr
 195 200 205
 Asn Tyr Asp Phe Leu Ser Pro Asn Tyr Thr Gly Gln Cys Pro Ser Gly
 210 215 220
 Ile Arg Ala Gln Asn Asp Gln Leu Gly Trp Leu Trp Gly Gln Ser Arg
 225 230 235 240
 Ala Leu Tyr Pro Ser Ile Tyr Met Pro Ala Val Leu Glu Gly Thr Gly
 245 250 255
 Lys Ser Gln Met Tyr Val Gln His Arg Val Ala Glu Ala Phe Arg Val
 260 265 270
 Ala Val Ala Ala Gly Asp Pro Asn Leu Pro Val Leu Pro Tyr Val Gln
 275 280 285
 Ile Phe Tyr Asp Thr Thr Asn His Phe Leu Pro Leu Asp Glu Leu Glu
 290 295 300
 His Ser Leu Gly Glu Ser Ala Ala Gln Gly Ala Ala Gly Val Val Leu
 305 310 315 320
 Trp Val Ser Trp Glu Asn Thr Arg Thr Lys Glu Ser Cys Gln Ala Ile
 325 330 335
 Lys Glu Tyr Met Asp Thr Thr Leu Gly Pro Phe Ile Leu Asn Val Thr
 340 345 350
 Ser Gly Ala Leu Leu Cys Ser Gln Ala Leu Cys Ser Gly His Gly Arg
 355 360 365
 Cys Val Arg Arg Thr Ser His Pro Lys Ala Leu Leu Leu Leu Asn Pro
 370 375 380
 Ala Ser Phe Ser Ile Gln Leu Thr Pro Gly Gly Gly Pro Leu Ser Leu
 385 390 395 400
 Arg Gly Ala Leu Ser Leu Glu Asp Gln Ala Gln Met Ala Val Glu Phe
 405 410 415
 Lys Cys Arg Cys Tyr Pro Gly Trp Gln Ala Pro Trp Cys Glu Arg Lys
 420 425 430
 Ser Met Trp
 435

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 21 amino acids
 - (B) TYPE: amino acid
 - (D) TOPOLOGY: linear

70

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ala Gly His Leu Leu Pro Ile Cys Ala Leu Phe Leu Thr Leu Leu
 1 5 10 15

Asp Met Ala Gln Gly
 20

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 435 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met Ala Gly His Leu Leu Pro Ile Cys Ala Leu Phe Leu Thr Leu Leu
 1 5 10 15

Asp Met Ala Gln Gly Phe Arg Gly Pro Leu Val Pro Asn Arg Pro Phe
 20 25 30

Thr Thr Val Trp Asn Ala Asn Thr Gln Trp Cys Leu Glu Arg His Gly
 35 40 45

Val Asp Val Asp Val Ser Val Phe Asp Val Val Ala Asn Pro Gly Gln
 50 55 60

Thr Phe Arg Gly Pro Asp Met Thr Ile Phe Tyr Ser Ser Gln Leu Gly
 65 70 75 80

Thr Tyr Pro Tyr Tyr Thr Pro Thr Gly Glu Pro Val Phe Gly Gly Leu
 85 90 95

Pro Gln Asn Ala Ser Leu Ile Ala His Leu Ala Arg Thr Phe Gln Asp
 100 105 110

Ile Leu Ala Ala Ile Pro Ala Pro Asp Phe Ser Gly Leu Ala Val Ile
 115 120 125

Asp Trp Glu Ala Trp Arg Pro Arg Trp Ala Phe Asn Trp Asp Thr Lys
 130 135 140

Asp Ile Tyr Arg Gln Arg Ser Arg Ala Leu Val Gln Ala Gln His Pro
 145 150 155 160

Asp Trp Pro Ala Pro Gln Val Glu Ala Val Ala Gln Asp Gln Phe Gln
 165 170 175

Gly Ala Ala Arg Ala Trp Met Ala Gly Thr Leu Gln Leu Gly Gly Ala
 180 185 190

Leu Arg Pro Arg Gly Leu Trp Gly Phe Tyr Gly Phe Pro Asp Cys Tyr
 195 200 205

Asn Tyr Asp Phe Leu Ser Pro Asn Tyr Thr Gly Gln Cys Pro Ser Gly
 210 215 220

Ile Arg Ala Gln Asn Asp Gln Leu Gly Trp Leu Trp Gly Gln Ser Arg
 225 230 235 240

Ala Leu Tyr Pro Ser Ile Tyr Met Pro Ala Val Leu Glu Gly Thr Gly
 245 250 255

Lys Ser Gln Met Tyr Val Gln His Arg Val Ala Glu Ala Phe Arg Val

	260		265		270
Ala Val	Ala Ala Gly Asp Pro	Asn Leu Pro Val	Leu Pro Tyr Val	Gln	
	275	280	285		
Ile Phe Tyr	Asp Thr Thr Asn His	Phe Leu Pro	Leu Asp Glu Leu	Glu	
	290	295	300		
His Ser Leu	Gly Glu Ser Ala Ala	Gln Gly Ala Ala	Gly Val Val	Leu	
	305	310	315	320	
Trp Val Ser	Trp Glu Asn Thr Arg Thr	Lys Glu Ser Cys	Gln Ala Ile		
	325	330	335		
Lys Glu Tyr	Met Asp Thr Thr Leu	Gly Pro Phe Ile	Leu Asn Val	Thr	
	340	345	350		
Ser Gly Ala	Leu Leu Cys Ser Gln	Ala Leu Cys Ser	Gly His Gly	Arg	
	355	360	365		
Cys Val Arg	Arg Thr Ser His Pro	Lys Ala Leu Leu	Leu Leu Asn	Pro	
	370	375	380		
Ala Ser Phe	Ser Ile Gln Leu Thr	Pro Gly Gly Gly	Pro Leu Ser	Leu	
	385	390	395	400	
Arg Gly Ala	Leu Ser Leu Glu Asp	Gln Ala Gln Met	Ala Val Glu	Phe	
	405	410	415		
Lys Cys Arg	Cys Tyr Pro Gly Trp	Gln Ala Pro Trp	Cys Glu Arg	Lys	
	420	425	430		
Ser Met Trp					
	435				

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2517 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TYCCTCCAGG AGTCTCTGGT GCAGCTGGGG TGGAACTCTG CCAGGCCCTG CTTAGGCCCC	60
CATCCTGGGG TCAGGAAATT TGGAGGATAA GGCCCTTCAG CCCAAGGTC AGCAGGGACG	120
AGCGGGCAGA CTGGCGGGTG TACAGGAGGG CTGGGTTGAC CTGYCCTTGG TCACTGAGGC	180
CATTGGATCT TCCTCCAGTG GCTGCCAGGA TTTCTGGTGG AAGAGACAGG AAGGCCTCCC	240
CCCCTTGGTC GGGTCAGCCT GGGGGCTGAG GGCCTGGCTG TCAGCCACTC TTCCCAGAAC	300
ATATGTCATG GCCTCAGTGG CTCATGGGGA AGCAGGGGTG GGCAGGCTTA GGCTAGAGCA	360
AGTCCTGTGG GAGATGGCAG AGGCCTGGTC TGAGAGGCAA CTCGGATGTG CCCTCCAGTG	420
GCCATGCTCC CCTCCATGCG TCTCCCTGTC CCTCTGGAG CCGTCAGGT CAATGTTTAA	480
CAGAAACCAG AGCAGCGGTG GATTAATGCG CAAGGGCTCA GCGCCCGAGC CCGTAGCAGT	540
GGGGGAATCG GAGACTTTGC AACCTGTTCT CAGCTCTGCC TCCCCTGGGC AGGTTGTCCT	600
CGACCAGTCC CGTGCCATGG CAGGCCACCT GCTTCCCATC TCGGCCCTCT TCCTGACCTT	660
ACTCGATATG GCCCAAGGCT TTAGGGGCCC CTGGGTACCC AACCGGECCT TCACCACCGT	720

CTGGAATGCA AACACCCAGT GGTGCCTGGA GAGGCACGGT GTGGACGTGG ATGTCAGTGT	780
CTTCGATGTG GTAGCCAACC CAGGGCAGAC CTTCCGCGGC CCTGACATGA CAATTTTCTA	840
TAGTCCCAG CTGGGCACCT ACCCCTACTA CACGCCCACT GGGGAGCCTG TGTITGGTGG	900
TCTGCCCCAG AATGCCAGCC TGATTGCCCA CCTGGCCCGC ACATTCCAGG ACATCCTGGC	960
TGCCATACCT GCTCCTGACT TCTCAGGGCT GGCAGTCATC GACTGGGAGG CATGGCGCCC	1020
ACGCTGGGCC TTCAACTGGG ACACCAAGGA CATTACCAGG CAGCGCTCAC GGGCACTGGT	1080
ACAGGCACAG CACCTGATT GGCAGCTCC TCAGGTGGAG GCAGTAGCCC AGGACCAGTT	1140
CCAGGGAGCT GCACGGGCTT GGATGGCAGG CACCCTCCAG CTGGGGGGGG CACTGCGTCC	1200
TGCGGGCCTC TGGGGCTTCT ATGGCTTCCC TGACTGCTAC AACTATGACT TTCTAAGCCC	1260
CAACTACACC GGCCAGTGCC CATCAGGCAT CCGTCCCAA AATGACCAGC TAGGGTGGCT	1320
GTGGGGCCAG AGCGGTGCC TCTATCCAG CATCTACATG CCCGCAGTGC TGGAGGGCAC	1380
AGGGAAGTCA CAGATGTATG TGCAACACCG TGTGGCCGAG GCATTCCGTG TGGCTGTGGC	1440
TGCTGGTGAC CCAATCTGC CGGTGCTGCC CTATGTCCAG ATCTTCTATG ACACGACAAA	1500
CCACTTCTG CCCCTGGATG AGCTGGAGCA CAGCCTGGGG GAGAGTGGG CCCAGGGGGC	1560
AGCTGGAGTG GTGCTCTGGG TGAGCTGGGA AAATACAAGA ACCAAGGAAT CATGTCAAGC	1620
CATCAAGSAG TATATGGACA CTACACTGGG GCCCTTCATC CTGAACGTGA CCAGTGGGGC	1680
CCTTCTCTGC AGTCAAGCCC TGTGCTCCGG CCATGGCCGC TGTGTCCGCC GCACCAGCCA	1740
CCCCAAGCC CTCCTCTCC TTAACCTGC CAGTTTCTCC ATCCAGCTCA GGCCTGGTGG	1800
TGGCCCCCTG AGCCTGCGGG GTGCCCTCTC ACTTGAAGAT CAGGCACAGA TGGCTGTGGA	1860
GTTCAAATGT CGATGCTACC CTGGCTGGCA GGCACCGTGG TGTGAGCGGA AGAGCATGTG	1920
GTGATTGGCC ACACACTGAG TTGCACATAT TGAGAACCTA ATGCACTCTG GGTCTGGCCA	1980
GGGCTTCTCT AAATACATGC ACAGTCATAC AAGTCATGGT CACAGTAAAG AGTACACTCA	2040
GCCACTGTCA CAGGCATATT CCTGCACAC ACATGCATAC TTACAGACTG GAATAGTGGC	2100
ATAAGGAGTT AGAACCACAG CAGACACCAT TCATTCTGC TCCATATGCA TCTACTGGC	2160
AAGGTCATAG ACAATTCTCT CAGAGACACT GAGCCAGTCT TTGAACGTGA GCAATCACAA	2220
AGGCTGACAT TCACTGAGTG CTAATCTTT GCCAATCCCC GTGCTAAGCG TTTTATGTGG	2280
ACTTATTCAT TCCTCACAAT GAGGCTATGA GGAACCTGAG TCACTCACAT TGAGAGTAAG	2340
CACGTTGCCC AAGGTTGCAC AGCAAGAAAA GGGAGAAGTT GAGATTCAAA CCCAGGCTGT	2400
CTAGCTCCGG GGGTACAGCC CTTGCACTCC TACTGAGTTC GTGTAACCA GCCCTGCACG	2460
ACCCCTGAAT CTGCTGAGAG GCACCACTCC AGCAAATAAA GCAGTCATGA TTTACTT	2517

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 20 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CAGGTTGTCC TGCACCACTC

20

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ATGTGCAACT CAGTGTGTGG C

21

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GTGCCATGGC AGGCCACC

18

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ATCACCACAT GCTCTCCGC

20

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 8 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Phe Arg Gly Pro Leu Leu Val Pro
1 5

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 4 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Cys Ala Ala Xaa

1

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 509 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met 1	Gly	Val	Leu	Lys	Phe	Lys	His	Ile	Phe	Phe	Arg	Ser	Phe	Val	Lys
Ser	Glu	Gly	Val	Ser	Gln	Ile	Val	Phe	Thr	Phe	Leu	Leu	Ile	Pro	Cys
Cys	Leu	Thr	Leu	Asn	Phe	Arg	Ala	Pro	Pro	Val	Ile	Pro	Asn	Val	Ala
Phe	Leu	Trp	Ala	Trp	Asn	Ala	Pro	Ser	Glu	Phe	Cys	Leu	Gly	Lys	Phe
Asp	Glu	Pro	Leu	Asp	Met	Ser	Leu	Phe	Ser	Phe	Ile	Gly	Ser	Pro	Arg
Ile	Asn	Ala	Thr	Gly	Gln	Gly	Val	Thr	Ile	Phe	Tyr	Val	Asp	Arg	Leu
Gly	Tyr	Tyr	Pro	Tyr	Ile	Asp	Ser	Ile	Thr	Gly	Val	Thr	Val	Asn	Gly
Gly	Ile	Pro	Gln	Lys	Ile	Ser	Leu	Gln	Asp	His	Leu	Asp	Lys	Ala	Lys
Lys	Asp	Ile	Thr	Phe	Tyr	Met	Pro	Val	Asp	Asn	Leu	Gly	Met	Ala	Val
Ile	Asp	Trp	Glu	Glu	Trp	Arg	Pro	Thr	Trp	Ala	Arg	Asn	Trp	Lys	Pro
Lys	Asp	Val	Tyr	Lys	Asn	Arg	Ser	Ile	Glu	Leu	Val	Gln	Gln	Gln	Asn
Val	Gln	Leu	Ser	Leu	Thr	Glu	Ala	Thr	Glu	Lys	Ala	Lys	Gln	Glu	Phe
Glu	Lys	Ala	Gly	Lys	Asp	Phe	Leu	Val	Glu	Thr	Ile	Lys	Leu	Gly	Lys
Leu	Leu	Arg	Pro	Asn	His	Leu	Trp	Gly	Tyr	Tyr	Leu	Phe	Pro	Asp	Cys
Tyr	Asn	His	His	Tyr	Lys	Lys	Pro	Gly	Tyr	Asn	Gly	Ser	Cys	Phe	Asn
Val	Glu	Ile	Lys	Arg	Asn	Asp	Asp	Leu	Ser	Trp	Leu	Trp	Asn	Glu	Ser
Thr	Ala	Leu	Tyr	Pro	Ser	Ile	Tyr	Leu	Asn	Thr	Gln	Gln	Ser	Pro	Val
Ala	Ala	Thr	Leu	Tyr	Val	Arg	Asn	Arg	Val	Arg	Glu	Ala	Ile	Arg	Val
Ser	Lys	Ile	Pro	Asp	Ala	Lys	Ser	Pro	Leu	Pro	Val	Phe	Ala	Tyr	Thr
Arg	Ile	Val	Phe	Thr	Asp	Gln	Val	Leu	Lys	Phe	Leu	Ser	Gln	Asp	Glu
Leu	Val	Tyr	Thr	Phe	Gly	Glu	Thr	Val	Ala	Leu	Gly	Ala	Ser	Gly	Ile
Val	Ile	Trp	Gly	Thr	Leu	Ser	Ile	Met	Arg	Ser	Met	Lys	Ser	Cys	Leu
Leu	Leu	Asp	Asn	Tyr	Met	Glu	Thr	Ile	Leu	Asn	Pro	Tyr	Ile	Ile	Asn
Val	Thr	Leu	Ala	Ala	Lys	Met	Cys	Ser	Gln	Val	Leu	Cys	Gln	Glu	Gln

75

370		375		380
Gly Val Cys Ile Arg	Lys Asn Trp Asn Ser Ser	Asp Tyr Leu His Leu		
385	390	395	400	
Asn Pro Asp Asn Phe Ala Ile Gln	Leu Glu Lys Gly Gly Lys Phe Thr			
	405	410	415	
Val Arg Gly Lys Pro Thr Leu Glu	Asp Leu Glu Gln Phe Ser Glu Lys			
	420	425	430	
Phe Tyr Cys Ser Cys Tyr Ser Thr	Leu Ser Cys Lys Glu Lys Ala Asp			
	435	440	445	
Val Lys Asp Thr Asp Ala Val Asp	Val Cys Ile Ala Asp Gly Val Cys			
	450	455	460	
Ile Asp Ala Phe Leu Lys Pro Pro Met	Glu Thr Glu Glu Pro Gln Ile			
465	470	475	480	
Phe Tyr Asn Ala Ser Pro Ser Thr	Leu Ser Ala Thr Met Phe Ile Val			
	485	490	495	
Ser Ile Leu Phe Leu Ile Ile Ser	Ser Val Ala Ser Leu			
	500	505		

What is claimed is:

CLAIMS

1. An antibody characterized by its ability to bind to native human plasma
5 hyaluronidase (hpHase).

2. An antibody that specifically binds to native human plasma
hyaluronidase (hpHase) produced by the process comprising the steps of:

(a) incubating a candidate antibody with a sample comprising native
10 hpHase, said incubating being for a time sufficient for formation of antibody-
hpHase complexes;

(b) contacting the sample with an insoluble support having anti-antibody and
detectably-labeled hyaluronic acid bound thereto for a time sufficient for formation
of anti-antibody-candidate antibody-hpHase complexes; and

15 (c) exposing the sample in contact with the support to an acidic pH of about
3.4 to 3.7, thereby allowing hpHase in the antibody-hpHase complex to degrade
the detectably labeled hyaluronic acid;

wherein samples associated with hyaluronic acid degradation comprise an
anti-hpHase antibody.

20

3. A hybridoma cell line 17E9 having ATCC accession number ATCC HB-
12213.

4. A hybridoma cell line 4D5 having ATCC accession number ATCC HB-
25 12214.

5. A method for screening candidate antibodies for binding to a native acid active hyaluronidase (aaHase), the method comprising the steps of:

(a) incubating a candidate antibody with a sample comprising native aaHase, said incubating being for a time sufficient for formation of antibody-
5 aaHase complexes;

(b) contacting the sample with an insoluble support having anti-antibody and detectably-labeled hyaluronic acid bound thereto for a time sufficient for formation of anti-antibody-candidate antibody-hpHase complexes; and

(c) exposing the sample in contact with the support to an acidic pH of about
10 3.4 to 3.7, thereby allowing hpHase in the antibody-aaHase complex to degrade the detectably labeled hyaluronic acid;

wherein samples associated with hyaluronic acid degradation comprise an anti-aaHase antibody.

15 6. A method of purifying human plasma hyaluronidase (hpHase) from a sample, the method comprising;

contacting a sample comprising hpHase with an anti-hpHase antibody, said contacting being for a time sufficient for formation of anti-hpHase antibody-hpHase complexes;

20 isolating hpHase from the complexes.

7. A device for immunopurification of native human plasma hyaluronidase (hpHase) comprising:

an insoluble support; and

25 an anti-hpHase antibody characterized by an ability to bind hpHase with a binding affinity of 10^7 l/mole or more.

8. An assay device for detection of hyaluronidase activity comprising:
an insoluble support;

biotinylated hyaluronic acid (bHA) covalently bound to the support, where
the biotinylated hyaluronic acid being the product of a one-step reaction of
5 hyaluronic acid, 1-ethyl-dimethylaminopropyl carbodiimide (EDC) and biotin
hydrazide.

9. A method for identifying a patient having or susceptible to a condition
associated with a LuCa-1 defect, the method comprising the steps of:

10 contacting a sample from the patient with an anti-hpHase antibody, the
sample being selected from the group consisting of tissue, blood, plasma, serum,
and urine, said contacting being for a time sufficient for formation of anti-hpHase
antibody-hpHase complexes;

detecting the amount of hpHase present in the sample; and

15 comparing the amount of hpHase detected in the sample with an amount
of hpHase in a control sample containing a known amount of hpHase correlated
with a normal level of hpHase;

wherein detection of an amount of hpHase in the patient sample that is less
than the amount of hpHase in the control sample is indicative of a LuCa-1 defect
20 in the patient.

10. Substantially purified human plasma hyaluronidase characterized by a
fatty acid moiety that is resistant to cleavage by phospholipase C, phospholipase
D, and N-glycosidase-F.

25

11. A method of purifying a native acid active hyaluronidase (aaHase)
from a sample, the method comprising the steps of:

(a) dissolving a sample suspected of containing an aaHase in a solution at
a temperature substantially less than room temperature, the solution comprising a
30 non-ionic detergent;

- (b) raising the temperature of the solution to a temperature substantially greater than room temperature, said raising resulting in the formation of a detergent-rich phase comprising aaHase and a detergent-poor phase; and
- (c) isolating aaHase from the detergent-rich phase.

5

12. An expression system for production of recombinant hpHase, the system comprising a transformed cell containing a nucleic acid construct comprising hpHase-encoding nucleic acid operably linked to a eukaryotic promoter.

10

13. A formulation for administration of human plasma hyaluronidase (hpHase) to a patient having a condition associated with a LuCa-1 gene defect comprising:

- a) a therapeutically effective amount of a substantially pure human plasma
- 15 hyaluronidase polypeptide; and
- b) a pharmaceutically acceptable carrier.

14. The use of human plasma hyaluronidase for the treatment or prevention of cancer associated with a LuCa-1 defect.

20

15. The use of a construct comprising a nucleotide sequence encoding a human plasma hyaluronidase polypeptide and a eukaryotic promoting sequence operably linked thereto for the treatment or prevention of cancer associated with a defective LuCa-1 gene.

25

16. An isolated recombinant human plasma hyaluronidase polypeptide.

17. A method of making a recombinant human plasma hyaluronidase polypeptide, the method comprising the steps of:

introducing into a host cell a nucleotide sequence encoding a human plasma hyaluronidase polypeptide;

5 culturing the host cell so as to allow for expression of the polypeptide from the introduced sequence; and

isolating human plasma hyaluronidase polypeptide.

1/9




FIG. 1




FIG. 2

2/9




FIG. 3




FIG. 4

4/9




FIG. 6




FIG. 7A




FIG. 7B




FIG. 8




FIG. 9

7/9

HSC-3 Orthotopic SCC Model
Peritumor Injections of Recombinant
Human Plasma HA'se




Fig. 10

8/9




Fig. 11

9/9




FIG. 12

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/18089

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/388.28, 413; 435/338, 7.95, 201, 814, 172.1; 436/518; 424/146.1, 94.62; 514/44; 935/14

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, REG, CA, BIOSIS, MEDLINE, CABA

search terms: hyaluronidase (and RN), human, plasma

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NATOWICZ et al. Human Serum Hyaluronidase: Characterization of a Clinical Assay. Clinica Chimica Acta. 09 February 1996, Vol. 245, No. 1, pages 1-6, see entire document.	5, 13-14
Y	AFIFY et al. Purification and Characterization of Human Serum Hyaluronidase. Arch. Biochem. Biophys. September 1993, Vol. 305, No. 2, pages 434-441, see entire document.	1-4, 7, 9-10, 13
A	Chem. Abst. Vol. 67, No. 7, issued 1967, page 2766, DE SALEGUI et al. 'The Existence of an Acid-Active Hyaluronidase in Serum.' Abstr. No. 29394. Arch. Biochem. Biophys. 1967, Vol. 120, No. 1, pages 60-67.	1-17

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

02 JANUARY 1998

Date of mailing of the international search report

03 FEB 1998

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JON P. WEBER, PH. D.

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US97/18089

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BOWNESS et al. Chromatographic Distinction Between Hyaluronidases from Human Serum and Ovine Testes. Biochim. Biophys. Acta. 08 January 1968, Vol. 151, No. 1, pages 288-290.	1-17
A	FISZER-SZAFARZ. B. Hyaluronidase Polymorphism Detected by Polyacrylamide Gel Electrophoresis. Application to Hyaluronidases from Bacteria, Slime Molds, Bee and Snake Venoms, Bovine Testes, Rat Liver Lysosomes and Human Serum. Anal. Biochem. 1984, Vol. 143, pages 76-81.	1-17
A	BOLLET et al. The Presence of Hyaluronidase in Various Mammalian Tissues. J. Biol. Chem. November 1963, Vol. 238, No. 11, pages 3522-3527.	1-17
X,P --- Y,P	FROST et al. Purification, Cloning, and Expression of Human Plasma Hyaluronidase. Biochem. Biophys. Res. Comm. 09 July 1997, Vol. 236, No. 1, pages 10-15, see entire document.	1-8, 10-13, 15-17 ----- 9, 14

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/18089

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☒

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/18089

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C12P 21/08; C07K 16/40; G01N 33/543; C12N 9/26, 15/52; A61K 38/46, 39/395, 31/70

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

530/388.28, 413; 435/338, 7.95, 201, 814, 172.1; 436/518; 424/146.1, 94.62; 514/44; 935/14

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1.

- I.
 - a) Claims 1-2, drawn to a first product, Ab to hpHase.
 - b) Claims 3-4, drawn to hybridoma cell lines that produce the Ab.
 - c) Claim 5, drawn to a first method of making, screening for Ab that bind to hpHase.
 - d) Claim 6, drawn to a first method of use, purifying hpHase with its Ab.
 - e) Claims 7-8, drawn to devices containing Ab to hpHase as key element.
- II. Claim 9, drawn to a second method of use, identifying a patient having LuCa-1 deficiency with an Ab to hpHase.
- III. Claims 10, 13 and 16, drawn to a second product, purified or recombinant hpHase and pharmaceutical compositions.
- IV. Claim 11, drawn to a first method of making the second product, hpHase, by non-recombinant means.
- V. Claim 12, drawn to a third product, an expression system for making recombinant hpHase.
- VI. Claim 14, drawn to a method of treating a patient with hpHase.
- VII. Claim 15, drawn to a method of treating a patient with DNA.
- VIII. Claim 17, drawn to a second method of making the second product, hpHase, by recombinant means.

The inventions listed as Groups I-VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Inventions Groups I, III and V are drawn to distinct products, an antibody, an enzyme and an expression system. While the antibody may bind to the enzyme specifically, so do ions, substrates and inhibitors. The system may be used to express the enzyme, but has no relationship to the antibody. The expression system produces the desired enzyme as well as many other products. None of the claimed products structurally resembles the others or functions in the same manner.

Inventions Groups I, IV and VIII are drawn to distinct methods of purifying the product of Group III, via antibody immunoaffinity, by conventional precipitation and chromatography, and by recombinant means. None of the processes shares any steps in common.

Group II is drawn to a second method of use of the product of Group I which is not immediately suggested by knowledge of the product. This method of use is unrelated to the method of Group I.

Groups VI and VII are distinct methods of treating a patient with an enzyme and with DNA. Neither process suggests the other.

Groups I and II are alternate methods of use of the antibody: to purify an enzyme, and to screen for a disease.

The hpHase is admitted to be known in the art at page 4, first full paragraph of the disclosure.